

## NATURAL HEALTH PRODUCT

# PSEUDOEPHEDRINE Children 6-11 years

This monograph is intended to serve as a guide to industry for the preparation of Product Licence Applications (PLAs) and labels for natural health product market authorization. It is not intended to be a comprehensive review of the medicinal ingredient.

#### Notes

- Text in parentheses is additional optional information which can be included on the label at the applicant's discretion.
- The solidus (/) indicates that the terms and/or statements are synonymous. Either term or statement may be selected by the applicant on the label.
- This monograph cannot be combined with any other monograph at Class II. Theses products may be submitted as a Class III application along with evidence to support their safety and efficacy.
- To mitigate the potential risk to the health of children, it is recommended that child resistant packaging/containers be used as described in Sections C.01.001(2) to (4), and subsection C.01.031(1) (a) (i) of the *Food and Drug Regulations* (JC 2024a).

#### **Compliance with Precursor Control Regulations:**

The *Precursor Control Regulations* (PCR) (JC 2024b) allows Canada to fulfill its international obligations with respect to the *United Nations Convention Against Illicit Traffic in Narcotic Drugs and Psychotropic Substances, 1988,* and provides a framework for the regulation of activities involving precursor chemicals which can be used in the production of illicit drugs and substances. Under the PCR, regulated chemicals are grouped into two classes: Class A and Class B. For Class A precursors such as pseudoephedrine and/or products containing them, persons wishing to be involved in activities such as importation, exportation, production, packaging, selling, and/or providing must first obtain a licence.

Further information regarding compliance with the PCR, including application forms and guidance documents pertaining to the application for a Class A precursor licence, is available at: http://www.healthcanada.gc.ca/precursors.

Date

April 25, 2025



### Proper name(s), Common name(s), Source information

Proper names(s)		Common names(s)	Source information
			Source ingredient(s)
٠	(1S,2S)-2-Methylamino-1-	Pseudoephedrine	<ul> <li>Pseudoephedrine</li> </ul>
	phenylpropan-1-ol		• Pseudoephedrine hydrochloride
•	(alphaS)-alpha-[(1S)-1-		• Pseudoephedrine sulfate
	(Methylamino)ethyl]benzenemethanol		
•	d-psi-Ephedrine		

References: Proper names: NIH 2024; RSC 2024; Common name: NIH 2024; RSC 2024; Source information: NIH 2024; RSC 2024.

### **Route of administration**

Oral

### **Dosage form(s)**

This monograph excludes foods or food-like dosage forms as indicated in the Compendium of Monographs Guidance Document.

Acceptable dosage forms for oral use are indicated in the dosage form drop-down list of the webbased Product Licence Application form for Compendial applications.

### Use(s) or Purpose(s)

- Used as a decongestant (for the common cold) (US FDA 2022; HC 2009; Williamson 2003; Mills and Bone 2000; Simons et al. 1996; Bright et al. 1981; Bye et al. 1980; Empey 1980).
- Temporarily relieves/For the temporary relief of cold/(and) flu symptoms such as nasal congestion/sinus congestion/stuffy nose/nasal stuffiness (US FDA 2022; HC 2009; Williamson 2003; Mills and Bone 2000; Simons et al. 1996; Bright et al. 1981; Bye et al. 1980; Empey et al. 1980).
- Temporarily relieves/For the temporary relief of nasal congestion/sinus congestion/stuffy nose/nasal stuffiness (due to hay fever/allergic rhinitis/the common cold) (US FDA 2022; HC 2009; Williamson 2003; Mills and Bone 2000; Simons et al. 1996; Bright et al. 1981; Bye et al. 1980; Empey et al. 1980).
- Nasal decongestant (US FDA 2022; HC 2009; Williamson 2003; Mills and Bone 2000; Simons et al. 1996; Bright et al. 1981; Bye et al. 1980; Empey 1980).

**Note:** The above usages can be combined on the product label (e.g., Nasal decongestant to temporarily relieve nasal congestion due to the common cold).



Dose(s)

## Subpopulation(s)

Children 6 to 11 years

## Quantity(ies)

30 – 120 milligrams pseudoephedrine per day; Not to exceed 30 milligrams per single dose (US FDA 2022; HC 2009; Simons et al. 1996; Dickerson et al. 1978).

### **Direction(s)** for use

- Give a single dose every 4-6 hours, up to 4 times per day (US FDA 2022; HC 2009).
- Do not give with any other cough and cold medications since harm may occur (HC 2009).

### **Duration(s) of use**

Ask a health care practitioner/health care provider/health care professional/doctor/physician for use beyond 7 days (HC 2009; US FDA 2004).

## **Risk information**

### Caution(s) and warning(s)

- Keep out of reach of children. In case of overdose, call a poison control centre or get medical help right away (US FDA 2022; HC 2009).
- Ask a health care practitioner/health care provider/health care professional/doctor/ physician if symptoms worsen or last for more than 7 days (US FDA 2022).
- Ask a health care practitioner/health care provider/health care professional/doctor/ physician if symptoms are accompanied by a high fever that lasts longer than 3 days, the production of thick yellow/green phlegm, rash or a persistent headache.
- Ask a health care practitioner/health care provider/health care professional/doctor/ physician before use if the child is taking any other medications including cough and cold or weight control medications or prescription drugs, or products which contain caffeine or ephedrine (Brinker 2010; HC 2009; Mills and Bone 2005; Greenway et al. 2004; Naik and Freudenberger 2004).
- Ask a health care practitioner/health care provider/health care professional/doctor/ physician before use if the child has asthma, diabetes, glaucoma, heart disease, high blood pressure, kidney disease, thyroid disease, or seizure disorders or any other serious medical condition (Ando et al. 2022; Ganesh et al. 2019; Brinker 2010; HC 2009; Eccles 2006; Jorge et al. 2006; Bensky et al. 2004; Williamson 2003; Blumenthal et al. 2000; Mills and Bone 2000; Simons et al. 1996; Lambert 1987).



### **Contraindication(s)**

**Do not use** in a child who is taking a monoamine oxidase inhibitor (MAOI) or for 2 weeks after stopping the MAOI drug (US FDA 2022; Brinker 2010; Eccles 2006; Mills and Bone 2005; Blumenthal et al. 2000).

#### Known adverse reaction(s)

- Stop use and ask a health care practitioner/health care provider/health care professional/ doctor/physician if the child experiences severe headache, confusion, visual changes/changes in vision, nervousness, dizziness, sleeplessness, difficulty or pain when urinating or allergic reactions (Ando et al. 2022; Chen 2022; US FDA 2022; Gewirtz et al. 2021; Hinduja 2020; Patel et al. 2020; Pilato 2020; Ducros 2012; Legriel et al. 2021; Mills and Bone 2005; Bensky et al. 2004; Blumenthal et al. 2000; Dickerson et al. 1978).
- Stop use and ask a health care practitioner/health care provider/health care professional/ doctor/physician if any new symptoms appear.

### **Non-medicinal ingredients**

Must be chosen from the current Natural Health Products Ingredients Database (NHPID) and must meet the limitations outlined in the database.

#### **Storage Conditions**

Must be established in accordance with the requirements described in the *Natural Health Products Regulations*.

#### All products (information for industry; not for labelling)

To be packaged in airtight container, protected from light (Ph.Eur. 2023; USP-NF 2023).

#### **Specifications**

- The finished product specifications must be established in accordance with the requirements described in the Natural and Non-prescription Health Product Directorate (NNHPD) Quality of Natural Health Products Guide.
- The medicinal ingredient must comply with the requirements outlined in the NHPID.



## **EXAMPLE OF PRODUCT FACTS:**

Consult the Guidance Document, Labelling of Natural Health Products for more details.

Product Facts			
Medicinal ingredient in each capsule			
Pseudoephedrine XX mg			
Uses			
• Temporarily relieve nasal congestion due to the common cold.			
Nasal decongestant.			
Warnings			
If applicable <sup>1</sup> :			
Allergens: food allergen, gluten (gluten source), sulphites			
Contains aspartame			
Do not use in a child who is taking a monoamine oxidase inhibitor (MAOI) or for 2 weeks after			
stopping the MAOI drug.			
Ask a health care practitioner before use • the child is taking any other medications including cough			
and cold or weight control medications or prescription drugs, or products which contain caffeine or			
ephedrine • the child has asthma, diabetes, glaucoma, heart disease, high blood pressure, kidney			
disease, thyroid disease, or seizure disorders or any other serious medical condition.			
Stop use and ask a health care practitioner if • symptoms worsen or last for more than 7 days •			
symptoms are accompanied by a high fever that lasts longer than 3 days, the production of thick			
yellow/green phlegm, rash or a persistent headache • the child experiences severe headache, confusion,			
visual changes, nervousness, dizziness, sleeplessness, difficulty or pain when urinating or allergic			
reactions • any new symptoms appear.			
Keep out of reach of children. In case of overdose, call a poison control centre or get medical help			
right away.			
Directions			
Children 6-11 years: • Give a single dose every 4-6 hours, up to 4 times per day • Do not give with			
any other cough and cold medications since harm may occur • Ask a health care practitioner for use			
beyond / days.			
Other information			
(Add any storage information)			
Non-medicinal ingredients			
List all NMIs			
Questions? (Call) 1-XXX-XXX-XXXX			

<sup>1</sup>This section can be removed from the table if the product contains no allergen or aspartame.

#### **References cited**

Ando Y, Ono Y, Sano A, Fujita N, Ono S. Posterior Reversible Encephalopathy Syndrome: A Review of the Literature. Intern Med. 2022;61(2):135-141.

Bensky D, Clavey S, Stoger E. Gamble A. Chinese Herbal Medicine: Materia Medica, 3RD Edition. Seattle (WA): Eastland Press; 2004.



Blumenthal M, Goldberg A, Brinkmann J, editors. Herbal Medicine: Expanded Commission E Monographs. Boston (MA): Integrative Medicine Communications; 2000.

Bright TP, Sandage Jr BW, Fletcher HP. Selected cardiac and metabolic responses to pseudoephedrine with exercise. The Journal of Clinical Pharmacology 1981;21:488-492.

Brinker F. Herb Contraindications and Drug Interactions, 4<sup>th</sup> edition. Sandy (OR): Eclectic Medical Publications; 2010.

Bye CE, Cooper J, Empey DW, Fowle ASE, Hughes DTD, Letley E, O'Grady J. Effects of pseudoephedrine and tripolidine, alone and in combination, on symptoms of the common cold. British Medical Journal 1980;281(6234):189-190.

Chen SP, Wang SJ. Pathophysiology of reversible cerebral vasoconstriction syndrome. J Biomed Sci. 2022;29(1):72.

Dickerson J, Perrier D, Mayersohn M, Bressler R. Dose tolerance and pharmacokinetic studies of L (+) pseudoephedrine capsules in man. European Journal of Clinical Pharmacology 1978;14:253-259.

Ducros A. Reversible cerebral vasoconstriction syndrome. Lancet Neurol. 2012;11(10):906-917.

Eccles R. 2006. Substitution of phenylephrine for pseudoephedrine as a nasal decongestant. An illogical way to control methamphetamine abuse. British Journal of Clinical Pharmacology 2006;63(1):10-14.

Empey DW, Young GA, Letley E, John GC, Smith P, McDonnell KA, Bagg LR, Hughes DTD. Dose-response study of the nasal decongestant and cardiovascular effects of pseudoephedrine. British Journal of Clinical Pharmacology 1980;9:351-358.

Ganesh K, Nair RR, Kurian G, Mathew A, Sreedharan S, Paul Z. Posterior Reversible Encephalopathy Syndrome in Kidney Disease. Kidney Int Rep. 2017;3(2):502-507.

Gewirtz AN, Gao V, Parauda SC, Robbins MS. Posterior Reversible Encephalopathy Syndrome. Current pain and headache reports 2021;5(3):19.

Greenway FJ, de Jonge L, Blanchard D, Frisard M, Smith, SR. Effect of a dietary herbal supplement containing caffeine and ephedra on weight, metabolic rate, and body composition. Obesity Research 2004;12(7):1152-1157.

HC 2009: Health Canada Guidance Document: Nonprescription oral paediatric cough and cold labelling standard. Health Products and Food Branch (HPFB). Health Canada. [Accessed 2025 April 7]. Available from: https://www.canada.ca/en/health-canada/services/drugs-health-products/drug-products/applications-submissions/guidance-documents/nonprescription-drugs-labelling-standards/nonprescription-oral-paediatric-cough-cold-labelling-standard.html

JC 2024a: Justice Canada. 2023. Food and Drug Regulations. Sections C.01.029, C.01.031 and C.01.031.2 (1). Cautionary Statements and Child Resistant Packages. Ottawa (ON): Department of

Pseudoephedrine - Children 6-11 years Page 6 of 9



Justice Canada. [Accessed 2025 April 7]. Available from: http://laws-lois.justice.gc.ca/eng/regulations/c.r.c.,\_c.\_870/page-97.html

JC 2024b: Justice Canada. 2023. Precursor Control Regulations. Part 1 – Class A Precursors. Ottawa (ON) Department of Justice Canada. [Accessed 2025 April 7]. Available from: http://laws-lois.justice.gc.ca/eng/regulations/SOR-2002-359/index.html

Jorge S, Lopes JA, De Almeida E, Martins Prata M. Encefalopatía posterior reversible [Posterior reversible encephalopathy syndrome (PRES) and chronic kidney disease]. Nefrologia. 2007;27(5):650 (Spanish).

Lambert MT. Paranoid psychoses after abuse of proprietary cold remedies. British Journal of Psychiatry 1987;151:548-550.

Legriel S, Schraub O, Azoulay E, et al. Determinants of recovery from severe posterior reversible encephalopathy syndrome. PLoS One 2012;7:e44534.

Mills S, Bone K. Principles and Practice of Phytotherapy. Toronto (ON): Churchill Livingstone; 2000.

Mills S, Bone K. The Essential Guide to Herbal Safety. St. Louis (MO): Elsevier Churchill Livingstone; 2005.

Naik SJ, Freudenberger RS. Ephedra–associated cardiomyopathy. The Annals of Pharmacotherapy 2004;38:400-403.

NIH 2024: National Institutes of Health. PubChem. Search terms: Pseudoephedrine, Pseudoephedrine Hydrochloride, Pseudoephedrine Sulfate. Bethesda (MD): National Library of Medicine, US Department of Health & Human Services. [Accessed 2025 April 7]. Available from: https://pubchem.ncbi.nlm.nih.gov/

Patel SD, Otite FO, Lima J, Khandelwal P, Ollenschleger M. Abstract WP207: Current Incidence, Epidemiological and Clinical Characteristics of Reversible Cerebral Vasoconstriction in the United States. Stroke 2020;51:AWP207,12.

Ph. Eur. 2023: European Pharmacopoeia Commission. 2007. European Pharmacopoeia, 11<sup>th</sup> edition. Strasbourg (FR): Directorate for the Quality of Medicines and HealthCare of the Council of Europe (EDQM).

Pilato F, Distefano M, Calandrelli R. Posterior Reversible Encephalopathy Syndrome and Reversible Cerebral Vasoconstriction Syndrome: Clinical and Radiological Considerations. Front Neurol, 2020;11:34.

RSC 2024: Royal Society of Chemistry: The Merck Index Online [Accessed 2025 April 7]. Available from: https://merckindex.rsc.org/

Simons FE, Gu X, Watson WT, Simons KJ. Pharmacokinetics of the orally administered decongestants pseudoephedrine and phenylpropanolamine in children. Journal of Pediatrics

Pseudoephedrine - Children 6-11 years Page 7 of 9



1996;129(5):729-734.

US FDA 2022: United States Food and Drug Administration. Over-the-Counter Monograph M012: Cold, Cough, Allergy, Bronchodilator and Antiasthmatic Drug Products for Over-the-Counter Human Use; Department of Health and Human Services, U.S. Food and Drug Administration. [Accessed 2025 March 31]. Available from: : https://dpsadmin.fda.gov/omuf/sites/omuf/files/primary-documents/2022-10/OTC%20Monograph\_M012-Cough%20Cold%20Allergy%20Bronchodilator%20and%20Antiasthmatic%20Drug%20Products%

20for%20OTC%20Human%20Use%2010.14.2022\_0.pdf

US FDA 2004: United States Food and Drug Administration. Final Rules Declaring Dietary Supplements Containing Ephedrine Alkaloids Adulterated Because They Present an Unreasonable Risk; Final Rule. Federal Register, Volume 69, Number 28, February 11, 2004, Rules and Regulations, Docket Number 1995N-0304. Rockville (MD): United States Department of Health and Human Services, U.S. Food and Drug Administration. [Accessed 2025 March 31]. Available from: https://www.gpo.gov/fdsys/pkg/FR-2004-02-11/pdf/04-2912.pdf

USP-NF 2023: United States Pharmacopeia and the National Formulary. Rockville (MD): United States Pharmacopeial Convention, Inc.; 2023.

Williamson EM. Potter's Herbal Cyclopaedia: The Authoritative Reference work on Plants with a Known Medical Use. Saffron Walden (GB): The C.W. Daniel Company Limited; 2003.

### **References reviewed**

Aaronson AL, Ehrlich NJ, Frankel DB, Gutman AA, Aaronson DW. Effective oral nasal decongestion. A double-blind crossover analysis. Annals of Allergy 1968;26(3):145-150.

Burke LM. Positive drug tests from supplements. Sportscience 2000;4(3):1-5.

Dawson JK, Earnshaw SM, Graham CS. Dangerous monoamine oxidase inhibitor interactions are still occurring in the 1990s. Journal of Accident and Emergency Medicine 1995;12:49-51.

Dingemanse J, Guentert T, Gieschke R, Stabl M. Modification of the cardiovascular effects of ephedrine by the reversible monoamine oxidase A-inhibitor moclobemide. Journal of Cardiovascular Pharmacology 1996;28(6):856-861.

Drew CD, Knight GT, Hughes DT, Bush M. Comparison of the effects of D-(-)-ephedrine and L-(+)-pseudoephedrine on the cardiovascular and respiratory systems in man. British Journal of Clinical Pharmacology 1978;6(3):221-225.

Elis J, Laurence DR, Mattie H, Prichard BNC. Modification by monoamine oxidase inhibitors of the effect of some sympathomimetics on blood pressure. British Medical Journal 1967;2:75-78.

Farel RM, Fanta CH. Drug therapy for asthma. New England Journal of Medicine 2009;360(24):2578-2579.



Gurley BJ, Gardner SF, White LM, Wang PL. Ephedrine pharmacokinetics after the ingestion of nutritional supplements containing Ephedra sinica (ma huang). Therapeutic Drug Monitoring 1998;20(4):439-445.

Haller CA, Jacob P 3rd, Benowitz NL. Enhanced stimulant and metabolic effects of combined ephedrine and caffeine. Clinical Pharmacology and Therapeutics 2004;75(4):259-273.

Herridge CF, A'Brook MF. Ephedrine psychosis. British Medical Journal 1968;2:160.

Hoffman D. Medical Herbalism: The Science and Practice of Herbal Medicine. Rochester (VT): Healing Arts Press; 2003.

Jacobs KM, Hirsch KA. Psychiatric complications of ma-huang. Psychosomatics 2000;41(1):58-62.

Laitinen LA, Empey DW, Bye C, Britton MG, McDonnell K, Hughes DTD. 1982. A comparison of the bronchodilator action of pseudoephedrine and ephedrine in patients with reversible airway obstruction. European Journal of Clinical Pharmacology 23:107-109.

MHRA 2009: MHRA Public Assessment Report - Controlling the risk of misuse of medicines containing pseudoephedrine and ephedrine. London (GB): Medicines and Healthcare products Regulatory Agency; 2009.

Mizoguchi H, Wilson A, Jerdack GR, Hull JD, Goodale M, Grender JM, Tyler BA. Efficacy of a single evening dose of syrup containing paracetamol, dextromethorphan hydrobromide, doxylamine succinate and ephedrine sulfate in subjects with multiple common cold symptoms. International Journal of Clinical Pharmacology and Therapeutics 2007;45(4):230-236.

Mortimer EA. Drug toxicity from breast milk? Pediatrics 1977;60(5):780-781.

Salem CB, Slim R, Denguezli M, Sriha B, Hmouda H, Bouraoui K. Pseudoephedrine induced acute generalized exanthematous pustulosis. The International Society of Dermatology 2008;47:418-419.

Slodki SJ, Montgomery CA. Clinical comparison of oxymetazoline and ephedrine in nasal congestion. Current Therapeutic Research, Clinical and Experimental 1965;7:19-22.

Theoharides TC. Sudden death of a healthy college student related to ephedrine toxicity from a ma huang-containing drink. Journal of Clinical Psychopharmacology 1997;17(5)437-439.

White LM, Gardner SF, Gurley BJ, Marx MA, Wang PL, Estes M. Pharmacokinetics and cardiovascular effects of ma huang (Ephedra sinica) in normotensive adults. The Journal of Clinical Pharmacology 1997;37:116-122.

Whitehouse AM, Duncan JM. Ephedrine psychosis rediscovered. British Journal of Psychiatry 1987;150:258-261.

Wingert WE, Mundy LA, Collins GL, Chmara ES. Possible role of pseudoephedrine and other over-the-counter cold medications in the deaths of very young children. Journal of Forensic Science 2007;52(2):487-490.

Pseudoephedrine - Children 6-11 years Page 9 of 9