

NATURAL HEALTH PRODUCT

COGNITIVE FUNCTION PRODUCTS

This monograph is intended to serve as a guide to industry for the preparation of Product Licence Applications (PLAs) and labels for natural health product market authorization. It is not intended to be a comprehensive review of the medicinal ingredients described within.

Notes

- Text in parentheses is additional optional information which can be included on the PLA and product label at the applicant's discretion.
- The solidus (/) indicates that the terms and/or the statements are synonymous. Either term or statement may be selected by the applicant.

Date March 12, 2021

Proper name(s), Common name(s), Source information

Table 1.1. Proper name(s), Common name(s), Source information for ingredients with nervine action

Proper name(s)	Common name(s)	Source information		
		Source material(s)	Part(s)	Preparation(s)
<i>Anemone pulsatilla</i>	<ul style="list-style-type: none"> • European pasqueflower • Pasqueflower • Pulsatilla 	<i>Anemone pulsatilla</i>	Herb top	Dried ¹
<i>Avena sativa</i>	<ul style="list-style-type: none"> • Common oat • Oat • Red oat • Side oat • Tree oat 	<i>Avena sativa</i>	Seed	Dried
<i>Nepeta cataria</i>	<ul style="list-style-type: none"> • Catmint • Catnep • Catnip • Catswort • Field balm 	<i>Nepeta cataria</i>	Herb top	Dried
<i>Matricaria chamomilla</i>	<ul style="list-style-type: none"> • Blue chamomile • Chamomile • Common chamomile • German chamomile • Hungarian chamomile • Matricaria • Scented chamomile • Scented mayweed 	<i>Matricaria chamomilla</i>	Flower	Dried

Proper name(s)	Common name(s)	Source information		
		Source material(s)	Part(s)	Preparation(s)
	<ul style="list-style-type: none"> • Sweet false chamomile • True chamomile • Wild chamomile 			
<i>Panax quinquefolius</i>	<ul style="list-style-type: none"> • American ginseng • Occidental ginseng • Sang • Shang • Xi yang shen 	<i>Panax quinquefolius</i>	Root	Dried
<i>Tilia cordata</i>	<ul style="list-style-type: none"> • Linden • Littleleaf linden • Small-leaf European linden • Small-leaf lime • Small-leaf lime tree • Small-leaf linden • Tilia 	<i>Tilia cordata</i>	Flower	Dried
<i>Tilia platyphyllos</i>	<ul style="list-style-type: none"> • Big-leaf linden • Broadleaf lime • Large-leaf lime • Large-leaf linden • Linden • Tilia 	<i>Tilia platyphyllos</i>	Flower	Dried
<i>Tilia x europaea</i>	<ul style="list-style-type: none"> • European Lime tree • European linden • Lime • Lime tree • Linden • Tilia 	<i>Tilia x europaea</i>	Flower	Dried
<i>Turnera diffusa</i>	Damiana	<i>Turnera diffusa</i>	<ul style="list-style-type: none"> • Leaf • Leaf and stem 	Dried
<i>Withania somnifera</i>	<ul style="list-style-type: none"> • Asgandh • Ashwagandha • Asvagandha • Winter-cherry • Withania 	<i>Withania somnifera</i>	Root	Dried

References: Proper names: NHPID 2019; Common names: NHPID 2019; Source information: Bone 2003; Hoffman 2003; Bradley 1992; Williamson et al. 1988; BHP 1983; Remington and Wood 1918; BPC 1911.

¹Do not use the fresh plant (Bone 2003; Hoffman 2003; Bradley 1992; BHP 1983).

Table 1.2. Proper name(s), Common name(s), Source information for ingredients with sedative action

Proper name(s)	Common name(s)	Source information		
		Source material(s)	Part(s)	Preparation(s)
<i>Eschscholzia californica</i>	California-poppy	<i>Eschscholzia californica</i>	Herb top	Dried
<i>Humulus lupulus</i>	<ul style="list-style-type: none"> • Bine • European hop • Hop • Hops 	<i>Humulus lupulus</i>	Strobile	Dried
<i>Hypericum perforatum</i>	<ul style="list-style-type: none"> • Goatweed • Hypericum • St. John's wort • St. John's-wort 	<i>Hypericum perforatum</i>	Herb top	Dried
<ul style="list-style-type: none"> • 5-Hydroxy-L-tryptophan • L-5-Hydroxytryptophan 	<ul style="list-style-type: none"> • L-5-HTP • L-5-Hydroxytryptophan 	<i>Griffonia simplicifolia</i> ¹	Seed	N/A
<i>Melissa officinalis</i>	<ul style="list-style-type: none"> • Balm • Bee balm • Lemon balm • Melissa 	<i>Melissa officinalis</i>	Herb top	Dried
<i>Passiflora incarnata</i>	<ul style="list-style-type: none"> • Apricot-vine • Maypop • Maypop passionflower • Passionflower 	<i>Passiflora incarnata</i>	Herb top	Dried
<i>Scutellaria lateriflora</i>	<ul style="list-style-type: none"> • Helmet flower • Hoodwort • Mad-dog scullcap • Quarter bonnet • Scullcap • Skullcap • Virginian skullcap 	<i>Scutellaria lateriflora</i>	Herb top	Dried
<i>Stachys officinalis</i>	<ul style="list-style-type: none"> • Bishop's-wort • Purple betony • Wood betony 	<i>Stachys officinalis</i>	Herb top	Dried
<i>Valeriana officinalis</i>	<ul style="list-style-type: none"> • Common valerian • Garden-heliotrope • Garden valerian • Valerian 	<i>Valeriana officinalis</i>	<ul style="list-style-type: none"> • Root • Root and rhizome 	Dried
<i>Withania somnifera</i>	<ul style="list-style-type: none"> • Asgandh • Ashwagandha • Asvagandha • Winter-cherry • Withania 	<i>Withania somnifera</i>	Root	Dried

References: Proper names: NHPID 2019; Common names: NHPID 2019; Source information: Hoffman 2003; Williamson et al. 1988.

¹Isolate

Table 1.3. Proper name(s), Common name(s), Source information for ingredients with cognitive support action

Proper name(s)	Common name(s)	Source information				
		Source ingredient(s)	Source material(s)	Organism group(s)	Part(s)	Preparation(s)
(2R)-2-(Acetyloxy)-3-carboxy-N,N,N-trimethyl-1-propanaminium inner salt	Acetylcarnitine	<ul style="list-style-type: none"> • Acetyl-L-carnitine arginate dihydrochloride¹ • Acetyl L-carnitine hydrochloride¹ 	N/A	N/A	N/A	N/A
<i>Bacopa monnieri</i>	<ul style="list-style-type: none"> • Bacopa • Moneywort • Water hyssop 	N/A	<i>Bacopa monnieri</i>	N/A	Herb top	Dried
<ul style="list-style-type: none"> • 2-[[[(2R)-2,3-Dihydroxypropoxy]hydroxyphosphinyl]oxy]-N,N,N-trimethylethanolaminium hydroxide inner salt • d-Choline hydroxide 2,3-dihydroxypropyl hydrogen phosphate inner salt • L-alpha-Glycerylphosphorylcholine 	Choline alfoscerate	Choline alfoscerate ¹	N/A	N/A	N/A	N/A
Cod liver oil	<ul style="list-style-type: none"> • Cod liver oil • Lecoris Aselli Oleum 	N/A	N/A	<ul style="list-style-type: none"> • <i>Melanogrammus aeglefinus</i> • <i>Arctogadus glacialis</i> 	Liver	N/A

Proper name(s)	Common name(s)	Source information				
		Source ingredient(s)	Source material(s)	Organism group(s)	Part(s)	Preparation(s)
				<ul style="list-style-type: none"> • <i>Gadus macrocephalus</i> • <i>Gadus morhua</i> • <i>Gadus ogac</i> • <i>Pollachius virens</i> 		
<i>Eleutherococcus senticosus</i>	<ul style="list-style-type: none"> • Ci wu jia • Eleuthero • Siberian ginseng 	N/A	<i>Eleutherococcus senticosus</i>	N/A	Root	Dried
Fish oil	Fish oil	N/A	N/A	<ul style="list-style-type: none"> • Ammodytidae • Carangidae • Clupeidae • Engraulida • Osmeridae • Salmonidae • Scombridae 	Whole	N/A
<i>Ginkgo biloba</i>	<ul style="list-style-type: none"> • Ginkgo • Ginkgo biloba • Maidenhair-tree • Baiguo • Icho 	N/A	<i>Ginkgo biloba</i>	N/A	Leaf	N/A
<i>Panax ginseng</i>	<ul style="list-style-type: none"> • Asian ginseng • Chinese ginseng • Hong shen • Korean ginseng • Korean red ginseng • Oriental ginseng • Panax ginseng 	N/A	<i>Panax ginseng</i>	N/A	Root Rootlet	Dried

Proper name(s)	Common name(s)	Source information				
		Source ingredient(s)	Source material(s)	Organism group(s)	Part(s)	Preparation(s)
	• Red ginseng • Ren shen					
Phosphatidylserine	Phosphatidylserine	N/A	<i>Helianthus annuus</i> ²	N/A	Seed	N/A
		Phosphatidylserine -enriched soy lecithin ¹	N/A	N/A	N/A	N/A
		Phosphatidylserine ¹	N/A	N/A	N/A	N/A
<i>Rhodiola rosea</i>	• Hong jing tian • King's-crown • Rhodiola • Roseroot	N/A	<i>Rhodiola rosea</i>	N/A	• Root • Root and rhizome	Dried
<i>Schizochytrium spp.</i>	Schizochytrium Oil	N/A	<i>Schizochytrium spp.</i>	N/A	Whole	N/A

References: Proper names: NHPID 2019; Common names: NHPID 2019; Source information: Yurko-Mauro et al. 2010; Calabrese et al. 2008; Bone 2003; De Jesus Moreno Moreno 2003; Parnetti et al. 2001; Barbagallo et al. 1994; Parnetti et al. 1993; Canal et al. 1991.

¹Synthetic

²Isolate

Table 1.4. Proper name(s), Common name(s), Source information for caffeine and ingredients containing caffeine

Proper name(s)	Common name(s)	Source information			
		Source ingredient(s)	Source material(s)	Part(s)	Preparation(s)
• 1,3,7-Trimethylxanthine • 3,7-Dihydro-1,3,7-trimethyl-1H-purine-2,6-dione	Caffeine	• Caffeine ¹ • Caffeine citrate ¹	N/A	N/A	N/A

Proper name(s)	Common name(s)	Source information			
		Source ingredient(s)	Source material(s)	Part(s)	Preparation(s)
<i>Ilex paraguariensis</i>	<ul style="list-style-type: none"> • Brazilian tea • Mate • Paraguayan tea • Paraguay tea • St Bartholomew's tea • Yerba mate 	N/A	<i>Ilex paraguariensis</i>	Leaf	Dried
<i>Paullinia cupana</i>	Guarana	N/A	<i>Paullinia cupana</i>	Seed	Dried

References: Proper names: NHPID 2019; Common names: NHPID 2019; Source information: EMA 2013; ESCOP 2009; Bradley 2006; Kennedy et al. 2004; Taylor 2003; Barnes et al. 2002; BHP 1983.

¹Synthetic

Table 1.5. Proper name(s), Common name(s), Source information for an ingredient with action on cognitive fatigue

Proper name(s)	Common name(s)	Source information	
		Source ingredient(s)	Preparation(s)
<ul style="list-style-type: none"> • (S)-alpha-Amino-4-hydroxybenzenepropanoic acid • L-Tyrosine 	<ul style="list-style-type: none"> • L-Tyrosine • Tyrosine 	<ul style="list-style-type: none"> • L-Tyrosine ethyl ester¹ • N-Acetyl-L-tyrosine¹ 	N/A

References: Proper names: NHPID 2019; Common names: NHPID 2019.

¹Synthetic

Table 1.6. Proper name(s), Common name(s), Source information for ingredients with action on memory

Proper name(s)	Common name(s)	Source information		
		Source material(s)	Part(s)	Preparation(s)
<i>Bacopa monnieri</i>	<ul style="list-style-type: none"> • Bacopa • Moneywort • Water hyssop 	<i>Bacopa monnieri</i>	<ul style="list-style-type: none"> • Whole plant • Herb top 	Dried
<i>Withania somnifera</i>	<ul style="list-style-type: none"> • Asgandh • Ashwagand • Asvagandha • Winter-cherry • Withania 	<i>Withania somnifera</i>	Root	Dried

References: Proper names: NHPID 2019; Common names: NHPID 2019; Source information: Bone 2003; API 2001.

Table 1.7. Proper name(s), Common name(s), Source information for ingredients with relaxation action

Proper name(s)	Common name(s)	Source information			
		Source ingredient(s)	Source material(s)	Part(s)	Preparation(s)
4-Aminobutanoic acid	<ul style="list-style-type: none"> • GABA • gamma-Aminobutyric acid 	N/A	<i>Lactobacillus hilgardii</i> ²	Whole cell for biosynthesis	N/A
		4-Aminobutanoic acid ¹	N/A	N/A	N/A
<ul style="list-style-type: none"> • L-Theanine • N-Ethyl-L-glutamine 	L-Theanine	N/A	<i>Camellia sinensis</i> ²	Leaf	N/A
		L-Theanine ¹	N/A	N/A	N/A

References: Proper names: NHPID 2019; Common names: NHPID 2019; Source information: Kanehira et al. 2011; Abdou et al. 2006; Yamakoshi et al. 2006.

¹Synthetic

²Isolate

Table 1.8. Proper name(s), Common name(s), Source information for ingredients with adaptogen action

Proper name(s)	Common name(s)	Source information		
		Source material(s)	Part(s)	Preparation(s)
<i>Asparagus racemosus</i>	<ul style="list-style-type: none"> • Shatamuli • Shatavari 	<i>Asparagus racemosus</i>	Root	Dried
<i>Astragalus membranaceus</i>	<ul style="list-style-type: none"> • Astragalus • Huangqi • Huang qi • Membranaceus milk vetch • Membranous milk-vetch • Milk vetch root • Mo jia huang qi 	<i>Astragalus membranaceus</i>	Root	Dried
<i>Eleutherococcus senticosus</i>	<ul style="list-style-type: none"> • Ci wu jia • Eleuthero • Siberian ginseng 	<i>Eleutherococcus senticosus</i>	<ul style="list-style-type: none"> • Root • Root and rhizome 	Dried
<i>Ganoderma lucidum</i>	<ul style="list-style-type: none"> • Ganoderma • Ling zhi • Reishi 	<i>Ganoderma lucidum</i>	<ul style="list-style-type: none"> • Cultured mycelium • Fruiting body 	Dried

Proper name(s)	Common name(s)	Source information		
		Source material(s)	Part(s)	Preparation(s)
			• Mycelium	
<i>Glycyrrhiza glabra</i>	• Guang guo gan cao • Licorice • Liquorice	<i>Glycyrrhiza glabra</i>	• Root • Root and stolon	Dried
<i>Ocimum tenuiflorum</i>	• Holy basil • Rama tulsi • Sacred basil • Tulsi	<i>Ocimum tenuiflorum</i>	Leaf	Dried
<i>Panax ginseng</i>	• Asian ginseng • Chinese ginseng • Hong shen • Korean ginseng • Korean red ginseng • Oriental ginseng • Panax ginseng • Red ginseng • Ren shen	<i>Panax ginseng</i>	• Root • Rootlet	Dried
<i>Panax quinquefolius</i>	• American ginseng • Occidental ginseng • Sang • Shang • Xi yang shen	<i>Panax quinquefolius</i>	Root	Dried
<i>Rhodiola rosea</i>	• Hong jing tian • King's-crown • Rhodiola • Roseroot	<i>Rhodiola rosea</i>	• Root • Root and rhizome	Dried
<i>Schisandra chinensis</i>	• Chinese Magnoliavine • Schisandra • Wu wei zi	<i>Schisandra chinensis</i>	Fruit	Dried
<i>Tinospora cordifolia</i>	Guduchi	<i>Tinospora cordifolia</i>	Stem	Dried
<i>Withania somnifera</i>	• Asgandh • Ashwagandha • Asvagandha • Winter-cherry • Withania	<i>Withania somnifera</i>	• Root • Whole plant	Dried

References: Proper names: NHPID 2019; Common names: NHPID 2019; Source information: Upton 2012; Winston and Kuhn 2008; Winston and Maimes 2007; WHO 2004; Bone 2003; Hoffman 2003; Thomsen 2002; Williamson 2002; Blumenthal et al. 2000; Upton 1999; WHO 1999; Bradley 1992; BHP 1983.

Route of administration

Oral

Dosage form(s)

This monograph excludes foods or food-like dosage forms as indicated in the Compendium of Monographs Guidance Document.

Acceptable dosage forms for the age category listed in this monograph and specified route of administration are indicated in the Compendium of Monographs Guidance Document.

Use(s) or Purpose(s) and Dose(s)

Note: Claims for traditional use must include the term “Herbal Medicine”, “Traditional Chinese Medicine”, or “Ayurveda”.

Subpopulation(s)

Adults 18 years and older

Quantity(ies)

Refer to Tables 2.1 to 2.8. below.

Table 2.1. Cognitive function uses or purposes and associated daily doses for ingredients with nervine action

Uses or Purposes ¹	Medicinal Ingredients	Methods of preparation	Doses/day ²
(Traditionally) used in Herbal Medicine to help relieve restlessness and/or nervousness (nervine/calmative)	<i>Anemone pulsatilla</i> (Pasqueflower)	Dry, Powder, Non-Standardized Ethanolic Extracts (Dry extract, Tincture, Fluid extract)	0.03-0.6 grams dried herb top, per day
		Non-Standardized Aqueous Extracts (Dry extract, Decoction, Infusion)	0.1-0.9 grams dried herb top, per day
	<i>Avena sativa</i> (Oat)	Dry, Powder, Non-Standardized Ethanolic Extracts (Dry extract, Tincture, Fluid extract)	0.6-20 grams dried seed, per day

Uses or Purposes ¹	Medicinal Ingredients	Methods of preparation	Doses/day ²
	<i>Matricaria chamomilla</i> (German chamomile)	Dry, Powder, Non-Standardized Extracts (Dry extract, Tincture, Fluid extract, Decoction, Infusion)	1.5-24 grams dried flower, per day
	<i>Nepeta cataria</i> (Catnip)	Dry, Powder, Non-Standardized Extracts (Dry extract, Tincture, Fluid extract, Decoction, Infusion)	1.2-12 grams dried herb top, per day
	<i>Panax quinquefolius</i> (American ginseng)	Dry, Powder, Non-Standardized Extracts (Dry extract, Tincture, Fluid extract, Decoction, Infusion)	0.5-12 grams dried root, per day
	<i>Tilia cordata</i> (Small-leaf linden)	Dry, Powder, Non-Standardized Extracts (Dry extract, Tincture, Fluid extract, Decoction, Infusion)	1.5-12 grams dried flower, per day
	<i>Tilia platyphyllos</i> (Large-leaf linden)	Dry, Powder, Non-Standardized Extracts (Dry extract, Tincture, Fluid extract, Decoction, Infusion)	1.5-12 grams dried flower, per day
	<i>Tilia x europaea</i> (European Linden)	Dry, Powder, Non-Standardized Extracts (Dry extract, Tincture, Fluid extract, Decoction, Infusion)	1.5-12 grams dried flower, per day
	<i>Turnera diffusa</i> (Damiana)	Dry, Powder, Non-Standardized Ethanolic Extracts (Dry extract, Tincture, Fluid extract)	1.5-4 grams dried leaf/leaf and stem, per day
Traditionally used in Ayurveda to balance aggravated Vata (nervine)	<i>Withania somnifera</i> (Ashwagandha)	Dry, Powder, Non-Standardized Extracts (Dry extract, Tincture, Fluid extract, Decoction, Infusion)	2-6 grams dried root, per day

¹References consulted for the uses or purposes: Bone 2003; Hoffman 2003; Ellingwood 1983; Felter and Lloyd 1983; Grieve 1971.

²References consulted for the doses: CNF 2012; Bone 2003; Hoffman 2003; Bradley 1992; Williamson et al. 1988; BHP 1983; Ellingwood 1983; Felter and Lloyd 1983; Remington and Wood 1918; BPC 1911.

Table 2.2. Cognitive function uses or purposes and associated daily doses for ingredients with calming sedative action

Uses or Purposes ¹	Medicinal Ingredients	Methods of preparation	Doses/day ²
(Traditionally) used in Herbal Medicine to help relieve restlessness and/or nervousness (calmative) AND/OR (Traditionally) used in Herbal Medicine as a sleep aid (during times of mental stress)	<i>Eschscholzia californica</i> (California poppy)	Dry, Powder, Non-Standardized Extracts (Dry extract, Tincture, Fluid extract, Decoction, Infusion)	0.2-3 grams dried herb top, per day
	<i>Humulus lupulus</i> (Hops)	Dry, Powder, Non-Standardized Extracts (Dry extract, Tincture, Fluid extract, Decoction, Infusion)	0.5-6 grams dried strobile, per day
	<i>Melissa officinalis</i> (Lemon balm)	Dry, Powder, Non-Standardized Extracts (Dry extract, Tincture, Fluid extract, Decoction, Infusion)	0.4-13.5 grams dried herb top, per day
	<i>Passiflora incarnata</i> (Passionflower)	Dry, Powder, Non-Standardized Ethanolic Extracts (Dry extract, Tincture, Fluid extract)	0.25-8 grams dried herb top, per day
		Non-Standardized Aqueous Extracts (Dry extract, Decoction, Infusion)	1-8 grams dried herb top, per day
	<i>Scutellaria lateriflora</i> (Skullcap)	Dry, Powder, Non-Standardized Ethanolic Extracts (Dry extract, Tincture, Fluid extract)	0.25-12 grams dried herb top, per day
		Non-Standardized Aqueous Extracts (Dry extract, Decoction, Infusion)	3-12 grams dried herb top, per day
	<i>Stachys officinalis</i> (Wood betony)	Dry, Powder, Non-Standardized Ethanolic Extracts (Dry extract, Tincture, Fluid extract)	2-4 grams dried herb top, per day
(Traditionally) used in Herbal Medicine to help relieve restlessness and/or nervousness (calmative)	<i>Hypericum perforatum</i> (St. John's wort)	Dry, Powder, Non-Standardized Extracts (Dry extract, Tincture, Fluid extract, Decoction, Infusion)	2-12.6 grams dried herb top, per day; Not to exceed 4.2 grams per single dose

Uses or Purposes ¹	Medicinal Ingredients	Methods of preparation	Doses/day ²
Used in Herbal Medicine to help relieve restlessness and/or nervousness (calmative) AND/OR (Used in Herbal Medicine to) help(s) relieve sleep disturbances associated with mood imbalance	<i>Hypericum perforatum</i> (St. John's wort)	Standardized Extracts (Dry extract)	600-1800 milligrams of extract, per day, standardized to 3-6% hyperforin and/or 0.12-0.28% hypericin; Not to exceed 600 milligrams of extract per single dose
Used as a sleep aid/to help promote sleep	L-5-Hydroxytryptophan (L-5-HTP)	Isolate	100-200 milligrams, per day
(Traditionally) used in Herbal Medicine to relieve restlessness and/or nervousness (calmative) AND/OR (Traditionally used in Herbal Medicine as a) sleep aid/(to) help(s) to promote sleep	<i>Valeriana officinalis</i> (Valerian)	Dry, Powder, Non-Standardized Extracts (Dry extract, Tincture, Fluid extract, Decoction, Infusion)	0.3-12 grams dried root/root and rhizome, per day; Not to exceed 3.6 grams per single dose
		Standardized Extracts (Dry extract, Tincture, Fluid extract, Decoction, Infusion)	0.3-12 grams of dried root or root and rhizome, per day; Not to exceed 3.6 grams per single dose and: 0.04-0.90% valerenic acid Or 0.10-0.90% sesquiterpenic acids
Traditionally used in Ayurveda as a sleep aid and/or to balance aggravated Vata (nervine)	<i>Withania somnifera</i> (Ashwagandha)	Dry, Powder, Non-Standardized Extracts (Dry extract, Tincture, Fluid extract, Decoction, Infusion)	2-6 grams dried root, per day

¹References consulted for the uses or purposes: Tobyn et al. 2010; Williamson et al. 1988.

²References consulted for the doses: Williamson et al. 1988.

Table 2.3. Cognitive function uses or purposes and associated daily doses for ingredients with cognitive support action

Uses or Purposes ¹	Medicinal Ingredients	Methods of preparation	Doses/day ²
Helps support cognitive health and/or brain function	Acetylcarnitine	N/A	1.5-4 grams, per day
	<i>Bacopa monnieri</i> (Bacopa)	Standardized Extracts (Dry extract)	300 milligrams of extract, per day, standardized to 40-55% bacosides
	Cod liver oil	Standardized fixed oil	18 years: 0.77-3.7 grams Cod liver oil, per day And 100-1,258 milligrams EPA + DHA including at least 100 milligrams DHA, per day And 138-2,800 micrograms RAE (vitamin A), per day And 1.15-23.12 micrograms vitamin D ₃ , per day 19 years and older: 0.77-4 grams Cod liver oil, per day And 100-1,360 milligrams EPA + DHA including at least 100 milligrams DHA, per day And 138-3,000 micrograms RAE (Vitamin A), per day And 1.15-25 micrograms Vitamin D ₃ , per day
	L-alpha-Glycerophosphoryl choline (Choline alfoscerate)	N/A	1.2 grams, per day

Uses or Purposes ¹	Medicinal Ingredients	Methods of preparation	Doses/day ²
Used in Herbal Medicine to help improve mental performance after periods of mental exertion	Fish oil	Standardized fixed oil	18 years: 100-2,500 milligrams EPA and DHA including at least 100 milligrams DHA, per day 19 years and older: 100-5,000 milligrams EPA and DHA including at least 100 milligrams DHA, per day
	Phosphatidylserine	N/A	300 milligrams, per day
	Schizochytrium oil	Standardized fixed oil	200-2,000 milligrams DHA, per day
	<i>Eleutherococcus senticosus</i> (Eleuthero)	Dry, Powder, Non-Standardized Extracts (Dry extract, Tincture, Fluid extract, Decoction, Infusion)	0.91-6 grams dried root, per day
Helps to enhance cognitive function and/or memory in adults	<i>Ginkgo biloba</i> (Ginkgo)	Standardized Extracts (Dry extract)	80-240 milligrams of extract, per day (50:1; quantity crude equivalent 4 - 12 g of dried leaves) and: 22-27% flavonoid glycosides And 5-7% terpene lactones
(Used in Herbal Medicine to) help(s) support cognitive function and/or reduce mental fatigue (in cases of mental stress) AND/OR Helps support cognition	<i>Panax ginseng</i> (Panax ginseng)	Dry, Powder, Non-Standardized Extracts (Dry extract, Tincture, Fluid Extract, Decoction, Infusion)	0.5-9 grams dried root/ rootlet, per day
		Standardized Extracts (Dry extract)	200-600 milligrams of extract, per day; standardized to 4-7% total ginsenosides; Not to exceed 9 grams of dried root/rootlet, per day
(Used in Herbal Medicine) (to) help(s) support cognitive function (such as	<i>Rhodiola rosea</i> (Rhodiola)	Tincture	1.2-1.8 grams of dried root/root and rhizome, per day

Uses or Purposes ¹	Medicinal Ingredients	Methods of preparation	Doses/day ²
mental focus and mental stamina)		Standardized Extracts (Dry extract)	144-680 milligrams of extract, per day; Not to exceed 200 milligrams per single dose and: 0.8 - 3% salidroside And/Or 1 - 6% rosavins

¹References consulted for the uses or purposes: Morgan and Stevens 2010; Yurko-Mauro et al. 2010; Calabrese et al. 2008; Malaguanera et al. 2008; De Jesus Moreno Moreno 2003; Parnetti et al. 2001; Stough et al. 2001; Thal et al. 2000; Barbagallo et al. 1994; Pettegrew et al. 1994; Parnetti et al. 1993; Sano et al. 1992; Canal et al. 1991; Spagnoli et al. 1991.

²References consulted for the doses: Jensen et al. 2010; Morgan and Stevens 2010; Quinn et al. 2010; Yurko-Mauro et al. 2010; Calabrese et al. 2008; Malaguanera et al. 2008; De Jesus Moreno Moreno 2003; Parnetti et al. 2001; Stough et al. 2001; Thal et al. 2000; Barbagallo et al. 1994; Pettegrew et al. 1994; Parnetti et al. 1993; Sano et al. 1992; Canal et al. 1991; Spagnoli et al. 1991.

Table 2.4. Cognitive function uses or purposes and associated daily doses for caffeine and ingredients containing caffeine

Uses or Purposes ¹	Medicinal Ingredients	Methods of preparation	Doses/day ²
Helps (temporarily) to promote alertness and wakefulness, and to enhance cognitive performance AND/OR Helps (temporarily) to relieve/reduce fatigue/tiredness AND/OR Helps (temporarily) to support/promote mental sharpness/alertness	1,3,7-Trimethylxanthine (Caffeine)	N/A	100-400 milligrams, per day; and 100-200 milligrams per single dose
	<i>Ilex paraguariensis</i> (Yerba mate)	Standardized Extracts (Dry extract, Tincture, Fluid extract, Decoction, Infusion)	Extract corresponding to a maximum Quantity Crude Equivalent of 6 grams dried leaf, per day and not exceeding 150 milligrams caffeine, per day; and providing 100-150 milligrams caffeine per single dose
	<i>Paullinia cupana</i> (Guarana)	Standardized Ethanolic Extracts (Dry extract, Tincture, Fluid extract)	Extract corresponding to a maximum Quantity Crude Equivalent of 3 grams dried seed, per day and not exceeding 200 milligrams caffeine per day; and providing 100-200 milligrams caffeine per single dose

Uses or Purposes ¹	Medicinal Ingredients	Methods of preparation	Doses/day ²
Used in Herbal Medicine to help temporarily promote alertness and wakefulness, and to enhance cognitive performance	<i>Ilex paraguariensis</i> (Yerba mate)	Standardized Ethanolic Extracts (Dry extract, Tincture, Fluid extract)	Extract corresponding to a Quantity Crude Equivalent of 3-6 grams dried leaf, per day and not exceeding 150 milligrams caffeine per day
		Standardized Aqueous Extracts (Dry extract, Decoction, Infusion)	Extract corresponding to a Quantity Crude Equivalent of 2-6 grams dried leaf, per day and not exceeding 150 milligrams caffeine per day
	<i>Paullinia cupana</i> (Guarana)	Standardized Ethanolic Extracts (Dry extract, Tincture, Fluid extract)	Extract corresponding to a Quantity Crude Equivalent of 1-3 grams dried seed, per day and not exceeding 200 milligrams caffeine per day

¹References consulted for the uses or purposes: EMA 2013; EMA 2010; ESCOP 2009; Bradley 2006; Christopher et al. 2005; Kennedy al. 2004; Taylor 2003; Barnes et al. 2002; Kamimori et al. 2000; Zwyghuizen- Doorenbos et al. 1990.

²References consulted for the doses: Health Canada 2018; EMA 2013; Health Canada 2012; EMA 2010; ESCOP 2009; Bradley 2006; Taylor 2003.

Table 2.5. Cognitive function uses or purposes and associated daily doses for ingredients with action on cognitive fatigue

Uses or Purposes ¹	Medicinal Ingredients	Methods of preparation	Doses/day ²
Helps to decrease cognitive fatigue due to physically stressful situations (e.g. extended wakefulness, exposure to cold, excessive noise)	L-Tyrosine	N/A	10-20 grams, per day; Not to exceed 10 grams per single dose

¹References consulted for the uses or purposes: Neri et al. 1995.

²References consulted for the doses: Mahoney et al. 2007.

Table 2.6. Cognitive function uses or purposes and associated daily doses for ingredients with action on memory

Uses or Purposes ¹	Medicinal Ingredients	Methods of preparation	Doses/day ²
Traditionally used in Ayurveda for memory enhancement	<i>Bacopa monnieri</i> (Bacopa)	Dry, Powder, Non-Standardized Ethanolic Extracts (Dry extract, Tincture, Fluid extract)	1-6.5 grams dried whole plant/herb top, per day
	<i>Withania somnifera</i> (Ashwagandha)	Dry, Powder, Non-Standardized Extracts (Dry extract, Tincture, Fluid extract, Decoction, Infusion)	2-6 grams dried root, per day
Helps support memory	<i>Bacopa monnieri</i> (Bacopa)	Standardized Extracts (Dry extract)	300 mg of extract, per day, standardized to 40-55% bacosides

¹References consulted for the uses or purposes: Barbhaya et al. 2008; Stough et al. 2008; Raghav et al. 2006; Murthy 2004; Roodenrys et al. 2002; API 2001.

²References consulted for the doses: Bone 2003; API 2001.

Table 2.7. Cognitive function uses or purposes and associated daily doses for ingredients with relaxation action

Uses or Purposes ¹	Medicinal Ingredients	Methods of preparation	Doses/day ²
Helps to temporarily promote relaxation	4-Aminobutanoic acid (GABA)	N/A	50-3,000 milligrams, per day; Not to exceed 750 milligrams per single dose
	L-Theanine	N/A	200-250 milligrams, per day

¹References consulted for the uses or purposes: Kanehira et al. 2011; Abdou et al. 2006.

²References consulted for the doses: Kanehira et al. 2011; Powers et al. 2007; Yamakoshi et al. 2006.

Table 2.8. Cognitive function uses or purposes and associated daily doses for ingredients with adaptogen action

Uses or Purposes ¹	Medicinal Ingredients	Methods of preparation	Doses/day ²
Used in Herbal Medicine as an Adaptogen to help increase energy and	<i>Asparagus racemosus</i> (Shatavari)	Dry, Powder, Non-Standardized Ethanolic Extracts (Dry extract, Tincture, Fluid extract)	2-6 grams dried root, per day

Uses or Purposes ¹	Medicinal Ingredients	Methods of preparation	Doses/day ²
resistance to stress (e.g. in case of mental and physical fatigue related to stress)	<i>Astragalus membranaceus</i> (Astragalus)	Dry, Powder, Non-Standardized Ethanolic Extracts (Dry extract, Tincture, Fluid extract)	2-4.8 grams dried root, per day
	<i>Eleutherococcus senticosus</i> (Eleuthero)	Dry, Powder, Non-Standardized Ethanolic Extracts (Dry extract, Tincture, Fluid extract)	1-6 grams dried root/ root and rhizome, per day
		Non-Standardized Aqueous Extracts (Dry extract, Decoction, Infusion)	2-3 grams dried root/ root and rhizome, per day
	<i>Ganoderma lucidum</i> (Reishi)	Dry, Powder, Non-Standardized Ethanolic Extracts (Dry extract, Tincture, Fluid extract)	1.5-6 grams dried cultured mycelium/ fruiting body/ mycelium, per day
		Standardized Ethanolic Extracts (Dry extract, Tincture, Fluid extract)	1.5-6 grams of dried cultured mycelium/ fruiting body/ mycelium, per day and not to exceed 40 % polysaccharides
		Decoction, Decoction concentrate	3-15 grams dried cultured mycelium/ fruiting body/ mycelium, per day
		Decoction Standardized, Decoction concentrate Standardized	3-15 grams dried cultured mycelium/ fruiting body/ mycelium, per day and not to exceed 40 % polysaccharides
	<i>Glycyrrhiza glabra</i> (Licorice)	Dry, Powder, Non-Standardized Ethanolic Extracts (Dry extract, Tincture, Fluid extract)	0.6-6 grams dried root/ root and stolon, per day
	<i>Ocimum tenuiflorum</i> (Holy basil)	Dry, Powder, Non-Standardized Ethanolic Extracts (Dry extract, Tincture, Fluid extract)	2-3 grams dried leaf, per day
		Non-Standardized Aqueous Extracts (Dry extract, Decoction, Infusion)	4.2-28 grams dried leaf, per day
	<i>Panax ginseng</i>	Dry, Powder, Non-	0.5-3 grams dried root/

Uses or Purposes ¹	Medicinal Ingredients	Methods of preparation	Doses/day ²
	(Panax ginseng)	Standardized Ethanolic Extracts (Dry extract, Tincture, Fluid extract)	rootlet, per day
		Non-Standardized Aqueous Extracts (Dry extract, Decoction, Infusion)	0.5-2 grams dried root/ rootlet, per day
	<i>Panax quinquefolius</i> (American ginseng)	Dry, Powder, Non-Standardized Ethanolic Extracts (Dry extract, Tincture, Fluid extract)	1-3 grams dried root, per day
	<i>Rhodiola rosea</i> (Rhodiola)	Non-Standardized Ethanolic Extracts (Dry extract)	144-400 milligrams of dry extract and a Quantity Crude Equivalent (QCE) of 216-2,000 milligrams of dried root/root and rhizome, per day; Not to exceed 200 milligrams of dry extract and a QCE of 1 gram of dried root/root and rhizome, per single dose
		Tincture	1.2-1.8 grams of dried root/root and rhizome, per day
		Standardized Extracts (Dry extract)	144-680 milligrams of extract, per day; Not to exceed 200 milligrams per single dose and: 0.8 - 3% salidroside And/Or 1 - 6% rosavins
	<i>Schisandra chinensis</i> (Schisandra)	Dry, Powder, Non-Standardized Ethanolic Extracts (Dry extract, Tincture, Fluid extract)	1.2-6 grams dried fruit, per day
		Non-Standardized Aqueous Extracts (Dry extract, Decoction, Infusion)	1.5-6 grams dried fruit, per day
	<i>Tinospora cordifolia</i> (Guduchi)	Dry, Powder, Non-Standardized Ethanolic Extracts (Dry extract,	0.9-3 grams dried stem, per day

Uses or Purposes ¹	Medicinal Ingredients	Methods of preparation	Doses/day ²
		Tincture, Fluid extract)	
		Decoction, Decoction concentrate	20-30 grams dried stem, per day
	<i>Withania somnifera</i> (Ashwagandha)	Dry, Powder, Non-Standardized Ethanolic Extracts (Dry extract, Tincture, Fluid extract)	2.5-6.5 grams dried root/whole plant, per day
(Used as an Adaptogen) (to) help(s) (to) temporarily relieve symptoms of stress (such as mental fatigue and sensation of weakness)	<i>Rhodiola rosea</i> (Rhodiola)	Non-Standardized Ethanolic Extracts (Dry extract)	144-400 milligrams of dry extract and a Quantity Crude Equivalent (QCE) of 216-2,000 milligrams of dried root/root and rhizome, per day; Not to exceed 200 milligrams of dry extract and a QCE of 1 gram of dried root/root and rhizome, per single dose
		Tincture	1.2-1.8 grams of dried root/root and rhizome, per day
		Standardized Extracts (Dry extract)	144-680 milligrams of extract, per day; Not to exceed 200 milligrams per single dose and: 0.8 - 3% salidroside And/Or 1 - 6% rosavins
Traditionally used in Ayurveda as Rasayana (rejuvenative tonic)	<i>Asparagus racemosus</i> (Shatavari)	Dry, Powder, Non-Standardized Ethanolic Extracts (Dry extract, Tincture, Fluid extract)	3-6 grams dried root, per day

¹References consulted for the uses or purposes: Upton 2012; Winston and Maimes 2007; Mills and Bone 2005; API 2004; WHO 2004; Bone 2003; Hoffman 2003; Thomsen 2002; Williamson 2002; Blumenthal et al. 2000; Upton 1999; WHO 1999; Bradley 1992.

²References consulted for the doses: Upton 2012; Winston and Maimes 2007; Mills and Bone 2005; API 2004; WHO 2004; Bone 2003; Hoffman 2003; Thomsen 2002; Williamson 2002; Blumenthal et al. 2000; Upton 1999; WHO 1999; Bradley 1992.

Direction(s) for use

See Table 4 below.

Combination rules and restrictions

All medicinal ingredients included in this monograph may be combined across all groups, with the following restrictions:

Use or Purpose Restrictions

- A use or purpose statement is only acceptable if at least one medicinal ingredient associated with that statement is present at a dose at or above the minimum daily dose listed in Table 2.
- Medicinal ingredients which do not meet the minimum daily dose for a use or purpose statement will be considered as acceptable complementary medicinal ingredients in product formulations.
- Traditional claims: When traditional claim(s) is/are made, each traditional claim must be preceded with the medicinal ingredient(s) that support(s) it, unless it is a single medicinal ingredient product.
- If products contain caffeine (Group 4), sedative/relaxation claims are not permitted.
- If products contain sedatives (Group 2), alertness claims are not permitted.

Rules for Caffeine

- The maximum amount of total caffeine permitted from all ingredients in the product is 400 mg/day, and 200 mg per single dose.
- The maximum daily amount of caffeine permitted for each medicinal ingredient must be respected. Please see Table 2 above.
- If the total amount of caffeine provided by the combination of ingredients in the product (e.g. Yerba mate and Guarana) is equal to or higher than 100 mg/day, risk statements from the NNHPD Caffeine Monograph are required. Please see Table 4 below.
- Additional synthetic caffeine added to Guarana and/or Yerba mate extracts must be represented as a separate medicinal ingredient.

Rules for sedatives

- When combining L-5-HTP, St. John's Wort, and/or Passionflower, an additional risk statement is required. Please see Table 4 below.
- If ingredients are combined within Group 2 (Ingredients with sedative action), the sum of the percentages of the maximum daily reference dose must not exceed 100%, for all ingredients contributing to support the claim. Please see Table 3 below.

Table 3. Combination Table to assess Safety of Group 2 Ingredients

Additive Indication	Group 2 (e.g. Traditionally used in Herbal Medicine as a sleep aid)		
Medicinal Ingredients	Maximum Daily Recommended Dose on PLA form	Maximum Daily Monograph Reference Dose	Percentage of the Maximum Daily Monograph Reference Dose (%)
<i>Eschscholzia californica</i> (California poppy)	0.5 g	3 g	0.5/3 = 16.67%
<i>Humulus lupulus</i> (Hops)	4 g	6 g	4/6 = 66.66%
<i>Hypericum perforatum</i> (St John's wort)	0.5 g	12.6 g	0.5/12.6 = 3.96%
Sum of Percentages			87.3%

The calculations are performed as follows:

- Percentage of the Maximum Daily Monograph Reference Dose = [(Maximum Daily Recommended Dose on PLA form) / (Maximum Daily Monograph Reference Dose)] x 100%

Safety assessment - Sum of Percentages:

- In the example above, the sum of Percentages of the Maximum Daily Monograph Reference Dose is 87.3%, which is less than 100%. Safety of the combination of ingredients is therefore supported.

Duration(s) of use

See Table 4 below.

Risk information

Consult Table 4 for appropriate statements related to each medicinal ingredient. The medicinal ingredients in Table 4 are associated with the required numbered statements below.

1. As per the respective NNHPD monograph.

Duration(s) of Use

2. Ayurvedic bacopa claim only: Use for at least 6 weeks to see beneficial effects.
3. *Products providing 300 mg or more GABA per day*: Consult a health care practitioner/health care provider/health care professional/doctor/physician for use beyond 4 weeks.

Caution(s) and warning(s)

4. Consult a health care practitioner/health care provider/health care professional/doctor/physician if symptoms persist or worsen.
5. Consult a health care practitioner/health care provider/health care professional/doctor/physician prior to use if you are pregnant or breastfeeding.
6. Consult a health care practitioner/health care provider/health care professional/doctor/physician prior to use if you are pregnant.
7. Consult a health care practitioner/health care provider/health care professional/doctor/physician prior to use if you are breastfeeding.
8. Stop use if you experience severe stomach and/or intestinal irritation.
9. Do not use if you have gastrointestinal irritation.
10. Consult a health care practitioner/health care provider/health care professional/doctor/physician prior to use if you have a urinary tract disorder and/or kidney disease.
11. Consult a health care practitioner/health care provider/health care professional/doctor/physician prior to use if you have liver disease, kidney disease and/or a seizure disorder.
12. Consult a health care practitioner/health care provider/health care professional/doctor/physician prior to use if you have high blood pressure.
13. *Products providing 300 mg or more GABA per day:* Do not use with alcohol.
14. Consult a health care practitioner/health care provider/health care professional/doctor/physician prior to use if you have glaucoma, and/or overactive bladder syndrome (detrusor instability).
15. Avoid taking with health products or foods that contain caffeine and/or increase blood pressure (e.g. medications, coffee, tea, colas, cocoa, guarana, mate, bitter orange extract, synephrine, octopamine, ephedra, ephedrine).
16. This product is not intended as a substitute for sleep.
17. *Combination of ingredients resulting in a total caffeine content of more than 300 mg per day:* Consult a health care practitioner/health care provider/health care professional/doctor/physician prior to use if you are of childbearing age.
18. Stop use and consult a health care practitioner/health care provider/health care professional/doctor/physician if you experience any of the following symptoms within a few hours after taking this product: changes in mental state such as restlessness or confusion, increased heart rate, enlarged pupils, loss of muscle coordination, sweating, shivering and/or gastrointestinal symptoms such as nausea, vomiting, diarrhea.

Contraindication(s)

19. Do not use this product if you are pregnant or breastfeeding.
20. Do not use this product if you are pregnant.

Known adverse reaction(s)

21. May cause digestive problems.

Table 4. Risk information, durations of use, and directions for use associated with medicinal ingredients

Medicinal Ingredients	Risk Information	Durations of Use	Directions for use
1,3,7-Trimethylxanthine (Caffeine)	1		1
4-Aminobutanoic acid (GABA)	4,5,13	3	
Acetyl-L-Carnitine (Acetylcarnitine)	5,11,21		
<i>Asparagus racemosus</i> (Shatavari)	6		
<i>Astragalus membranaceus</i> (Astragalus)	1		
<i>Avena sativa</i> (Oat)	4		
<i>Bacopa monnieri</i> (Bacopa)	5,21	2	
Cod liver oil	1		
<i>Eleutherococcus senticosus</i> (Eleuthero)	1	1	
<i>Eschscholzia californica</i> (California poppy)	1		
Fish oil	1		
<i>Ganoderma lucidum</i> (Reishi)	1		1
<i>Ginkgo biloba</i> (Ginkgo)	1	1	
<i>Glycyrrhiza glabra</i> (Licorice)	1	1	
<i>Humulus lupulus</i> (Hops)	1		
<i>Hypericum perforatum</i> (St. John's wort)	1	1	
<i>Ilex paraguariensis</i> (Yerba mate)	5,12		
L-5-Hydroxytryptophan (L-5-HTP)	1		1
L-alpha-Glycerophosphorylcholine (Choline alfoscerate)	5		
L-Theanine	1		
L-Tyrosine	1	1	1
<i>Matricaria chamomilla</i> (German chamomile)	1		
<i>Melissa officinalis</i> (Lemon balm)	1		
<i>Nepeta cataria</i> (Catnip)	1		
<i>Ocimum tenuiflorum</i> (Holy basil)	1		
<i>Panax ginseng</i> (Asian ginseng)	1	1	
<i>Panax quinquefolius</i> (American ginseng)	1		
<i>Passiflora incarnata</i> (Passionflower)	1		
<i>Paullinia cupana</i> (Guarana)	5,12		
Phosphatidylserine	1		
<i>Anemone pulsatilla</i> (Pasqueflower)	4,8,9,10,19		
<i>Rhodiola rosea</i> (Rhodiola)	1		
<i>Schisandra chinensis</i> (Schisandra)	7,20		
<i>Scutellaria lateriflora</i> (Skullcap)	1		
<i>Stachys officinalis</i> (Wood Betony)	4,5		
<i>Tilia cordata</i> (Small-leaf linden)	1		
<i>Tilia platyphyllos</i> (Large-leaf linden)	1		
<i>Tilia x europaea</i> (European linden)	1		

<i>Tinospora cordifolia</i> (Guduchi)	5		
<i>Turnera diffusa</i> (Damiana)	4,7,20		
<i>Valeriana officinalis</i> (Valerian)	1		
<i>Withania somnifera</i> (Ashwagandha)	1		
Combination of ingredients resulting in a total caffeine content of 100 mg/day or more (e.g. Yerba mate and Guarana)	as per the caffeine monograph		
Combination of two or more of the following ingredients: L-5-HTP, St John's Wort, and/or Passionflower	18		

References consulted: EMA 2013; Gardner and McGuffin 2013; Fiebich et al. 2011; AMR 2010; Morgan and Stevens 2010; Okun et al. 2010; ESCOP 2009; Calabrese et al. 2008; CPS 2008; Cornelis and El-Sohemy 2007; Powers et al. 2007; Bain et al. 2006; Bradley 2006; Bui et al. 2006; Mills et al. 2006; Shils et al. 2006; Turner et al. 2006; Bouchard et al. 2005; Chandrasekaran et al. 2005; Haller et al. 2005; Noordzij et al. 2005; FDA 2004; Nathan et al. 2004; Bone 2003; Erner et al. 2003; Avisar et al. 2002; Barnes et al. 2002; Berardi et al. 2002; Thomsen 2002; Brinker 2001; Nathan et al. 2001; Stough et al. 2001; Arya et al. 2000; Vahedi et al. 2000; Jee et al. 1999; Mester et al. 1995; Bradley 1992; Zimmerman 1992; Creighton and Stanton 1990; Rai et al. 1990; FDA 1988; Jefferson 1988.

Non-medicinal ingredients

Must be chosen from the current Natural Health Products Ingredients Database (NHPID) and must meet the limitations outlined in that database.

Storage conditions

Products containing Cod liver oil, Fish oil or Schizochytrium oil

- Refrigerate after opening (except when encapsulated) (Senanayake and Fichtali 2006; Wille and Gonus 1989).
- Store in airtight container, protected from light (Ph.Eur. 2012; USP 35 2012).

Specifications

- The finished product specifications must be established in accordance with the requirements described in the Natural and Non-prescription Health Products Directorate (NNHPD) Quality of Natural Health Products Guide.
- The medicinal ingredient must comply with the requirements outlined in the NHPID.
- For the following ingredients, please see the respective NNHPD monograph for additional quality requirements: Astragalus, American ginseng, Cod liver oil, Eleuthero, Fish oil, German chamomile-Oral, Gingko biloba, Hops, Licorice, Panax ginseng, St John's wort, L-Tyrosine, L-Theanine, Valerian.

References cited

Abdou AM, Higashiguchi S, Horie K, Kim M, Hatta H, Yokogoshi H. Relaxation and immunity enhancement effects of gamma-aminobutyric acid (GABA) administration in humans. *Biofactors* 2006; 26(3):201-8.

AMR 2010: Alternative Medicine Review (AMR). Acetyl-L-Carnitine Monograph. Alternative Medicine Review 2010; 15(1): 76-83.

API 2001: The Ayurvedic Pharmacopoeia of India, Part I, Volume II. Delhi (IN): The Controller of Publications; 2001.

API 2004: The Ayurvedic Pharmacopoeia of India, Part 1, Volume IV. Delhi (IN): The Controller of Publications; 2004.

Arya LA, Myers DL, Jackson ND. Dietary caffeine intake and the risk for detrusor instability: a case-control study. *Obstetrics and Gynecology* 2000; 96(1):85-89.

Avisar R, Avisar E, Weinberger D. Effect of coffee consumption on intraocular pressure. *The Annals of Pharmacotherapy* 2002; 36(6):992-995.

Bain MA, Faull R, Fornasini G. Accumulation of trimethylamine and trimethylamine-N-oxide in end-stage renal disease patients undergoing haemodialysis. *Molecular Genetics and Metabolism* 2004; 81:263-272.

Barbagallo SG, Barbagallo M, Giordano M, Meli M, Panzarasa R. Alpha-glycerophosphocholine in mental recovery of cerebral ischemic attacks. An Italian multicenter clinical trial. *Annals of the New York Academy of Science* 1994; 717: 253-269.

Barbhaya C, Desai R, Saxena V, Pravina K, Wasim P, Geetharani P, Allan J, Venkateshwarlu K, Amit A. Efficacy and tolerability of bacomin® on memory improvement in elderly participants - A double blind placebo controlled study. *Journal of Pharmacology and Toxicology* 2008; 3(6): 425-434.

Barnes J, Anderson LA, Philipson JD. *Herbal Medicines: A Guide for Healthcare Professionals*, 2nd edition. London (GB): The Pharmaceutical Press; 2002.

Berardi RR, DeSimone EM, Newton GD, Oszko MA, Popovich NG, Rollins CJ, Shimp LA, Tietze KJ, editors. *Handbook of Non-prescription Drugs: An Interactive Approach to Self-Care*, 13th edition. Washington (DC): American Pharmaceutical Association; 2002.

BHP 1983: *British Herbal Pharmacopoeia*. Cowling (GB): British Herbal Medical Association; 1983.

Blumenthal M, Goldberg A, Brinckmann J. *Herbal Medicine: Expanded Commission E Monographs*. American Botanical Council; 2000.

Bone K. A clinical guide to blending liquid herbs: Herbal formulations for the individual patient. St. Louis (MI): Churchill Livingstone; 2003.

Bouchard NC, Howland MA, Greller HA, Hoffman RS, Nelson LS. Ischemic stroke associated with use of an ephedra-free dietary supplement containing synephrine. Mayo Clinic Proceeding 2005; 80(4):541-545.

BPC 1911: British Pharmaceutical Codex. Council of the Pharmaceutical Society of Great Britain; 1911.

Bradley PR, editor. British Herbal Compendium: A Handbook of Scientific Information on Widely Used Plant Drugs, Volume 1. Bournemouth (GB): British Herbal Medicine Association; 1992.

Bradley PR, editor. British Herbal Compendium: A Handbook of Scientific Information on Widely Used Plant Drugs, Volume 2. Bournemouth (GB): British Herbal Medicine Association; 2006.

Brinker F. Herb Contraindications and Drug Interactions, 3rd edition. Sandy (OR): Eclectic Medical Publications; 2001.

Brinker F. 2010. Online Updates and Additions to Herb Contraindications and Drug Interactions, 3rd edition. Sandy (OR): Eclectic Medical Publications; 2010. [Accessed 2014-03-04]. Available from: <http://www.eclecticherb.com/emp/updatesHCDI.html>.

Bui LT, Nguyen DT, Ambrose PJ. Blood pressure and heart rate effects following a single dose of bitter orange. The Annals of Pharmacotherapy 2006; 40(1):53-57.

Calabrese C, Gregory WL, Leo M, Kraemer D, Bone K, and Oken B. Effects of a standardized Bacopa monnieri extract on cognitive performance, anxiety, and depression in the elderly: A randomized, double-blind, placebo-controlled trial. The Journal of Alternative and Complementary Medicine 2008; 14(6): 707-713.

Canal N, Franceschi M, Alberoni M, Castiglioni C, De Moliner P, Longoni A. Effects of L-alpha-glyceryl-phosphorylcholine on amnesia caused by scopolamine. International Journal of Clinical Pharmacology. Therapy and Toxicology 1991; 29(3): 103-107.

Chandrasekaran S, Rochtchina E, Mitchell P. Effects of caffeine on intraocular pressure: the Blue Mountains Eye Study. Journal of Glaucoma 2005;14(6):504-507.

Christopher G, Sutherland D, Smith A. Effects of caffeine in non-withdrawn volunteers. Human Psychopharmacology 2005;20(1):47-53.

CNF 2012: Canadian Nutrient File (CNF). Nutrition & Healthy Eating, Food and Nutrition, Health Canada. [Accessed 2014-03-26]. Available from: <http://webprod3.hc-sc.gc.ca/cnf-fce/index-eng.jsp>

Cornelis MC, El-Sohemy A. Coffee, caffeine, and coronary heart disease. *Current Opinion in Lipidology* 2007;18(1):13-19.

CPS 2008: Compendium of Pharmaceuticals and Specialties: The Canadian Drug Reference for Health Professionals. Ottawa (ON): Canadian Pharmacists Association; 2008.

Creighton SM, Stanton SL. Caffeine: does it affect your bladder? *British Journal of Urology* 1990;66(6):613-614.

De Jesus Moreno Moreno M. Cognitive improvement in mild to moderate Alzheimer's dementia after treatment with the acetylcholine precursor choline alfoscerate: A multicenter, double-blind, randomized, placebo-controlled trial. *Clinical Therapeutics* 2003; 25(1): 178-193.

Ellingwood F. 1983. American Materia Medica, Therapeutics and Pharmacognosy. Sandy (OR): Eclectic Medical Publications [Reprint of 1919 original].

EMA 2010: European Medicines Agency. Community herbal monograph on *Ilex paraguariensis* St. Hilaire, folium. London (UK): Committee on Herbal Medicinal Products (HMPC); 2009. [Accessed 29 May 2017]. Available from: http://www.ema.europa.eu/docs/en_GB/document_library/Herbal_-_Community_herbal_monograph/2010/01/WC500059234.pdf.

EMA 2013: European Medicines Agency. Community herbal monograph on *Paullinia cupana* Kunth ex H.B.K. var. *sorbilis* (Mart.) Ducke, semen. London (UK): Committee on Herbal Medicinal Products (HMPC); 2013. [Accessed 29 May 2017]. Available from: http://www.ema.europa.eu/docs/en_GB/document_library/Herbal_-_Community_herbal_monograph/2013/03/WC500140768.pdf.

Erner RA, Meglatheray SB, Van Decker WA, Gallagher RM. Serotonin Syndrome and Other Serotonergic Disorders. *Pain Medicine* 2003; 4(1): 63-74.

ESCOP 2009: European Scientific Cooperative on Phytotherapy. ESCOP Monographs: The Scientific Foundation for Herbal Medicinal Products, 2nd edition, Supplement 2009. Exeter (GB): European Scientific Cooperative on Phytotherapy in collaboration with Thieme.

FDA 1988: Food and Drug Administration. 21 CFR Part 340. Stimulant drug products for over-the-counter human use; final monograph; final rule. Washington (DC): U.S. Food and Drug Administration, Department of Health and Human Services; 1988. [Accessed 2007-05-30]. Available from: http://ecfr.gpoaccess.gov/cgi/t/text/text-idx?c=ecfr&tpl=/ecfrbrowse/Title21/21cfr340_main_02.tpl.

FDA 2004: Food and Drug Administration. Sales of supplements containing ephedrine alkaloids (ephedra) prohibited. Washington (DC): Food and Drug Administration, US Department of Health and Human Services; 2004. [Accessed 2007-04-13]. Available from: <http://www.fda.gov/oc/initiatives/ephedra/february2004/>.

Felter HW, Lloyd JU. King's American Dispensatory, Volume 1, 18th edition. Sandy (OR): Eclectic Medical Publications; 1983 [Reprint of 1898 original].

Fiebich BL, Knorle R, Appel K, Kammler T, Weiss G. Pharmacological studies in an herbal drug combination of St. John's Wort (*Hypericum perforatum*) and passion flower (*Passiflora incarnata*): In vitro and in vivo evidence of synergy between *Hypericum* and *Passiflora* in antidepressant pharmacological models 2011; 82: 474-480.

Gardner Z and McGuffin M, editors. 2013. American Herbal Products Association's Botanical Safety Handbook Second Edition. Boca Raton (FL): CRC Press.

Grieve M. 1971. A Modern Herbal, Volume 1. New York (NY): Dover Publications [Reprint of 1931 Harcourt, Brace & Company publication].

Haller CA, Benowitz NL, Jacob P. Hemodynamic effects of ephedra-free weight-loss supplements in humans. The American Journal of Medicine 2005;118(9):998-1003.

Health Canada 2018. Caffeine monograph. Natural and Non-Prescription Health Products Directorate. Ottawa (ON). [Accessed 2019-02-14]. Available from: <http://webprod.hc-sc.gc.ca/nhpid-bdipsn/atReq.do?atid=caffeine.caffeine&lang=eng>

Health Canada 2012. Caffeine in foods. Health Canada. [Accessed 2019-02-14]. Available from: <https://www.canada.ca/en/health-canada/services/food-nutrition/food-safety/food-additives/caffeine-foods/foods.html>

Hoffmann D. Medical Herbalism: The Science and Practice of Herbal Medicine. Rochester (VT): Healing Arts Press; 2003.

Jee SH, He J, Whelton PK, Suh II, Klag MJ. The effect of chronic coffee drinking on blood pressure: a meta-analysis of controlled clinical trials. Hypertension 1999;33(2):647-652.

Jefferson JW. Lithium tremor and caffeine intake: two cases of drinking less and shaking more. Journal of Clinical Psychiatry 1988;49(2):72-73.

Jensen CL Voigt RG, Llorente AM, Peters SU, Prager TC, Zou YL, Rozelle JC, Turchich, MR, Frayley JK, Anerson RE, Heird WC. Effects of early maternal docosahexaenoic acid intake on neuropsychological status and visual acuity at five years of age of breast-fed term infants. The Journal of Pediatrics 2010; 125:900-905.

Kamimori H, Penetar DM, Headley DB, Thorne DR, Otterstetter R, Belenky G. Effect of three caffeine doses on plasma catecholamines and alertness during prolonged wakefulness. European Journal of Clinical Pharmacology 2000;56(8):537-544.

Kanehira T, Nakamura Y, Nakamura K, Horie K, Horie N, Furugori K, Sauchi Y, Yokogoshi H. Relieving Occupational Fatigue by Consumption of a Beverage Containing gamma-Amino Butyric Acid. Journal of Nutritional Science and Vitaminology 2011; 57(1):9-15.

Kennedy DO, Haskell CF, Wesnes KA, Scholey AB (2004). Improved cognitive performance in human volunteers following administration of guarana (*Paullinia cupana*) extract: comparison and interaction with Panaxginseng. *Pharmacology, Biochemistry, and Behaviour* 2004; 79(3):401-411.

Mahoney CR, Castellani J, Kramer FM, Young A, Lieberman HR. Tyrosine supplementation mitigates working memory decrements during cold exposure. *Physiology and Behavior* 2007;92(4):575-582.

Malaguarnera M, Gargate, MP, Cristaldi E, Colonna V, Messano M, Koverech A, Neri S, Vacante M, Cammalleri L, Motta, M. Acetyl L-carnitine (ALC) treatment in elderly patients with fatigue. *Archives of Gerontology and Geriatrics* 2008; 46: 181-190.

Mester R, Toren P, Mizrahi I, Wolmer L, Karni N, Weizman A. Caffeine withdrawal increases lithium blood levels. *Biological Psychiatry* 1995;37(5):348-350.

Mills E, Dugoua JJ, Perri D, Koren G. *Herbal Medicines in Pregnancy and Lactation*. Boca Raton (FL):Taylor and Francis Medical; 2006.

Mills S and Bone K. *The Essential Guide to Herbal Safety*. St. Louis (MO): Elsevier Churchill Livingstone; 2005.

Morgan and Stevens. Does Bacopa monnieri improve memory performance in older persons? Results of a randomized, placebo-controlled, double-blind trial. *The Journal of Alternative and Complementary Medicine* 2010; 16 (7): 753-759.

Murthy KRS. *Bhavaprakasha of Bhavmisra*, Volume 1. Varanasi (IND): Chowkhamba Krishnadas Academy; 2004.

Nathan PJ, Clarke, J, Lloyd J, Hutchison CW, Downey L, Stough C. The acute effects of an extract of Bacopa monniera (Brahmi) on cognitive function in healthy normal subjects. *Human Psychopharmacology* 2001; 16:345-351.

Nathan PJ, Tanner S, Lloyd J, Harrison B, Curran, L, Oliver C, Stough C. Effects of a combined extract of Ginkgo biloba and Bacopa monniera on cognitive function in healthy humans. *Human Psychopharmacology* 2004; 19(2):91 -96.

Neri DF, Wiegmann D, Stanny RR, Shappell SA, McCardie A, McKay DL. The effect of tyrosine on cognitive performance during extended wakefulness. *Aviation, space, and environmental medicine* 1995;66(4):313-319.

NHPID 2019. Natural Health Products Ingredients Database. [Accessed 2019-02-14]. Available from: <http://webprod.hc-sc.gc.ca/nhpidsn/search-rechercheReq.do>

Noordzij M, Uiterwaal CS, Arends LR, Kok FJ, Grobbee DE, Geleijnse JM. Blood pressure response to chronic intake of coffee and caffeine: a meta-analysis of randomized controlled trials. *Journal of Hypertension* 2005;23(5):921-928.

Okun MS, Boothby LA, Bartfield RB, Doering PL. GHB: an important pharmacologic and clinical update. *Journal of Pharmacy & Pharmaceutical Sciences: A Publication of the Canadian Society for Pharmaceutical Sciences, Société canadienne des sciences pharmaceutiques*. 2001; 4(2):167-175.

Parnetti L, Abate G, Bartorelli L, Cucinotta D, Cuzzupoli M, Maggioni M, Villardita C, Senin U. Multicentre study of l- α -glyceryl-phosphorylcholine vs ST200 among patients with probable senile dementia of Alzheimer's type. *Drugs and Aging* 1993; 3(2): 159-164.

Parnetti L, Amenta F, and Gallai, V. Choline alphoscerate in cognitive decline and in acute cerebrovascular disease: an analysis of published clinical data. *Mechanisms of Ageing and Development* 2001; 122:2041-2055.

Pettegrew JA, Klunk WE, Panchalingam K, Kanfer JN, McClure RJ. Clinical and neurochemical effects of acetyl-l-carnitine in Alzheimer's disease. *Neurobiology of Aging* 1994; 16(1):1-4.

Ph.Eur. 2012: European Pharmacopoeia, 8th edition. Strasbourg (FR): Directorate for the Quality of Medicines and HealthCare of the Council of Europe (EDQM); 2012.

Powers ME, Yarrow JF, McCoy SC, Borst SE. Growth Hormone Isoform Responses to GABA Ingestion at Rest and after Exercise. *Medicine & Science in Sports & Exercise* 2007;40(1):104- 110.

Quinn JF, Raman R, Thomas RG, Yurko-Mauro K, Nelson EB, Van Dyck C, Galvin JE, Emond J, Jack CR, Weiner M, Shinto L, Aisen PS. Docosahexaenoic acid supplementation and cognitive decline in Alzheimer's disease. *Journal of the American Medical Association* 2010;304(17): 1903-1911.

Raghav S, Singh H, Dalal P, Srivastava J, Asthana O. Randomized controlled trial of standardized *Bacopa monniera* extract in age-associated memory impairment. *Indian Journal of Psychiatry* 2006; 48(4): 238-242.

Rai G, Wright G, Scott L, Beston B, Rest J, Exton-Smith AN. Double-blind, placebo controlled study of acetyl-L-carnitine in patients with Alzheimer's dementia. *Current Medical Research Opinion* 1990; 11:638-647.

Remington JP and Woods HC, editors. *The Dispensatory of the United States of America*, 20th edition; 1918.

Roodenrys S, Booth D, Bulzomi S, Phipps A, Micallef C, Smoker J. Chronic effects of Brahmi (*Bacopa monnieri*) on human memory. *Neuropsychopharmacology* 2002; 27(2):279- 81.

Sano M, Bell K, Cote L, Dooneief G, Lawton A, Legler L, Marder K, Naini A, Stern Y, Mayeux R. Double blind parallel design pilot study of acetyl levocarnitine in patients with Alzheimer's disease. *Archive of Neurology* 1992; 49 (11): 1137-1141.

Senanayake SPJN, Fichtali J. Single-cell oils as sources of nutraceutical and specialty lipids: processing technologies and applications. In: Shahidi F, editor. *Neutraceutical and speciality lipids and their co-products*. Boca Raton (FL): Taylor and Francis Group; 2006.

Shils ME, Olson JA, Shike M, Ross AC, editors. *Modern Nutrition in Health and Disease*, 10th Cognitive Function Products Page 32 of 35

edition. Philadelphia (PA): Lippincott Williams and Wilkins; 2006.

Spagnoli A, Lucca U, Menasce, Bandera L, Cizza G, Forloni G, Tettamanti M, Frattura L, Tiraboschi P, Comelli M, et al. Long-term acetyl-L-carnitine treatment in Alzheimer's disease. *Neurology* 1991; 41(11):1726-1732.

Stough C, Downey L, Lloyd J, Silber B, Redman S, Hutchison C, Wesnes K, Nathan P. Examining the nootropic effects of a special extract of Bacopa monniera on human cognitive functioning: 90 day double-blind placebo-controlled randomized trial. *Phytotherapy Research* 2008; 22(12):1629-34.

Stough C, Lloyd J, Clarke J, Downey L, Hutchison CW, Rodgers T, Nathan PJ. The chronic effects of an extract of Bacopa monniera (Brahmi) on cognitive function in healthy human subjects. *Psychopharmacology* 2001; 156(4): 481-484

Taylor L. Technical data report for Yerba mate: Reprinted from *Herbal Secrets of the Rainforest*, 2nd Edition. Sage Press Inc.; 2003.

Thal LJ, Calvani M, Amato A, Carta AA. 1-year controlled trial of acetyl-L-carnitine in early-onset AD. *Neurology* 2000; 55: 805-810.

Thomsen M. Shatavari-Asparagus racemosus, Herbal Monograph, Phytomedicine. NSW Australia; 2002.

Tobyn G, Denham A, Whitelegg M. *The Western Herbal Tradition: 2000 years of medicinal plant knowledge*. Churchill-Livingstone; 2010.

Turner EH, Loftis JM, Blackwell AD. Serotonin a la carte: supplementation with the serotonin precursor 5-hydroxytryptophan. *Pharmacology & Therapeutics* 2006;109(3):325-338.

Upton R, editor. *American Herbal Pharmacopoeia and Therapeutic Compendium: American Ginseng Root (Panax quinquefolius L.)-Standards of Analysis, Quality Control, and Therapeutics*. Santa Cruz (CA): American Herbal Pharmacopeia; 2012.

Upton R, editor. *American Herbal Pharmacopoeia and Therapeutic Compendium: Schisandra Berry (Schisandra chinensis)-Analytical, Quality Control, and Therapeutic Monograph*. Santa Cruz (CA): American Herbal Pharmacopoeia; 1999.

USP 35 2012: United States Pharmacopeia and the National Formulary (USP 35 - NF 30). Rockville (MD): The United States Pharmacopeial Convention; 2012.

Vahedi K, Domingo V, Amarenco P, Bousser MG. Ischaemic stroke in a sportsman who consumed MaHuang extract and creatine monohydrate for body building. *Journal of Neurology, Neurosurgery and Psychiatry* 2000;68(1):112-113.

Wille HJ, Gonos P. Preparation of Fish Oil for Dietary Applications. In: Galli C, Simopolous AP, editors. *Dietary ω3 and ω6 Fatty Acids. Biological Effects and Nutritional Essentiality*. New York (NY): Plenum Press; 1989.

Williamson EM, editor. Major Herbs of Ayurveda. Edinburgh (GB): Churchill Livingstone; 2002

Williamson EM, Evans FJ, Wren RC. Potter's New Cyclopaedia of Botanical Drugs and Preparations. Saffron Walden (GB): C.W. Daniel Company Limited; 1988.

Winston D, Maimes S. Adaptogens: Herbs for strength, stamina and stress relief. Rochester (VT): Healing Arts Press; 2007.

Winston D and Kuhn MA. Herbal Therapy and Supplements: A Scientific and Traditional Approach. Philadelphia (PA): Wolters Kluwer Health and Lippincott Williams & Wilkins; 2008.

WHO: World Health Organization. WHO monographs on selected medicinal plants 2004; Volume 2.

WHO: World Health Organization. WHO monographs on selected medicinal plants 1999; Volume 1.

Yamakoshi J, Shiojo R, Nakagawa S, Izui N, Ogihaara T. Hypotensive Effects and Safety of Less-salt Soy Sauce Containing gamma-aminobutyric acid (GABA) on High-Normal Blood Pressure and Mild Hypertensive Subjects. *Yakuri To Chirya* 2006; 34:691-709 [English Translation].

Yurko-Mauro K, McCarthy D, Rom D, Nelson EB, Ryan AS, Blackwell A, Salem N, Stedman M. Beneficial effects of docosahexaenoic acid on cognition in age-related cognitive decline, *Alzheimer's and Dementia* 2010; 6:1-9.

Zimmerman DR. Zimmerman's Complete Guide to Non-prescription Drugs, 2nd edition. Detroit (MI): Gale Research Inc.; 1992.

Zwick H, Brush J, Iacullo CM, Connelly E, Gregory WL, Soumyanath A, Buresh R. The effect of Echinacea purpurea, astragalus membranaceus and Glycyrrhiza glabra on CD25 Expression in Humans: A Pilot Study. *Phytotherapy Research* 2007; 21(11): 1109-1112.

Zwyghuizen-Doorenbos A, Roehrs TA, Lipschutz L, Timms V, Roth T. Effects of caffeine on alertness. *Psychopharmacology* 1990; 100(1):36-39.

References reviewed

Blumenthal M, Busse W, Goldberg A, Gruenwald, J, Hall T, Riggins C, Rister R. The complete German Commission E monographs. Austin, TX: American Botanical Council; 1998.

Brinker F. The Toxicology of Botanical Medicines. Sandy (OR): Eclectic Medical Publications; 2000.

Calvani, M., Carta, A., Benedetti, N., Iannuccelli, M., Caruso, G. Action of acetyl-L-carnitine in neurodegeneration and Alzheimer's disease. *Aging and Cellular Defense Mechanisms* 1992;

663:483-486.

Gogte VVM. Ayurvedic Pharmacology and Therapeutic Uses of Medicinal Plants. Mumbai (IN): Bharatiya Vidya Bhavan; 2000.

Rege NN, Thatte UM, and Dahanukar SA. Adaptogenic properties of sixrasayana herbs used in Ayurvedic medicine. *Phytotherapy Research* 1999; 13(4):275-291.

Sayre, L. A Manual of Organic Materia Medica and Pharmacognosy. P. Blakiston's Son & Company; 1917.

Upton R, editor. American Herbal Pharmacopoeia and Therapeutic Compendium: Astragalus Root (*Astragalus membranaceus* & *Astragalus membranaceus* var. *mongolicus*)-Analytical, Quality Control, and Therapeutic Monograph. Santa Cruz (CA): American Herbal Pharmacopoeia; 1999.

Wilson L. Review of adaptogenic mechanisms: *Eleutherococcus senticosus*, *Panaxginseng*, *Rhodiola rosea*, *Schisandra chinensis* and *Withania somnifera*. *Australian Journal of Medical Herbalism* 2007; 19(3): 126-131.

Zwick H, Brush J, Iacullo CM, Connelly E, Gregory WL, Soumyanath A, Buresh R. The effect of *Echinacea purpurea*, *astragalus membranaceus* and *Glycyrrhiza glabra* on CD25 Expression in Humans: A Pilot Study. *Phytotherapy Research* 2007; 21(11): 1109-1112.