

NATURAL HEALTH PRODUCT

CAFFEINE

This monograph is intended to serve as a guide to industry for the preparation of Product Licence Applications (PLAs) and labels for natural health product market authorization. It is not intended to be a comprehensive review of the medicinal ingredient.

Notes

- ▶ Text in parentheses is additional optional information which can be included on the PLA and product label at the applicant's discretion.
- ▶ The solidus (/) indicates that the terms and/or statements are synonymous. Either term or statement may be selected by the applicant.

Date

October 30, 2018

Proper name(s), Common name(s), Source material(s)

Table 1. Proper name(s), Common name(s), Source material(s)

Proper name(s)	Common name(s)	Source material(s)
		Common name(s)
<ul style="list-style-type: none"> ▶ 1,3,7-Trimethylxanthine ▶ 3,7-Dihydro-1,3,7-trimethyl-1H-purine-2,6-dione 	Caffeine	<ul style="list-style-type: none"> ▶ Caffeine ▶ Caffeine citrate

References: Proper names: USP 30 2007; Common name: USP 30 2007, IOM 2003; Source materials: Ashihara and Suzuki 2004, Zajac et al. 2003, Gennaro 2000.

Route of administration

Oral (Higdon and Frei 2006)

Dosage form(s)

This monograph does not apply to caffeine added to food. Caffeine is regulated as a food additive when it is added to foods, including beverages, chewing gum, and bars. Questions about using caffeine in food can be sent to the Food Directorate (food-aliment@hc-sc.gc.ca).

Acceptable dosage forms for the age category listed in this monograph and specified route of administration are indicated in the Compendium of Monographs Guidance Document.

Use(s) or Purpose(s)

- ▶ Helps (temporarily) to promote alertness and wakefulness, and to enhance cognitive performance (Christopher et al. 2005; Kamimori et al. 2000; Zwyghuizen-Doorenbos et al. 1990).
- ▶ Helps (temporarily) to support/promote mental sharpness/alertness (Christopher et al. 2005; Kamimori et al. 2000; Zwyghuizen-Doorenbos et al. 1990).
- ▶ Helps (temporarily) to relieve fatigue, to promote endurance, and to enhance motor performance (Philip et al. 2006; Doherty and Smith 2005; Smith et al. 2005).
- ▶ Helps (temporarily) to enhance (physical) energy (Philip et al. 2006; Doherty and Smith 2005; Smith et al. 2005).
- ▶ Helps (temporarily) to reduce tiredness and fatigue (Philip et al. 2006; Doherty and Smith 2005; Smith et al. 2005).
- ▶ Used (temporarily) as a mild diuretic (Shirley et al. 2002; Neuhäuser-Berthold et al. 1997).

Dose(s)

Subpopulation(s)

Adults 18 years and older

Quantity(ies)

Mild diuretic

100 - 200 milligrams caffeine, per single dose; Not to exceed 800 milligrams, per day (Shirley et al. 2002; IOM 2001; Neuhäuser-Berthold et al. 1997).

Other uses

100 - 200 milligrams caffeine, per single dose; Not to exceed 1000 milligrams, per day (Sawynok 1995; FDA 1988; Greden 1974).

Direction(s) for use

Products providing more than 200 mg caffeine, per day (i.e. to be taken in divided doses)

Wait 3 to 4 hours between each dose.

Duration(s) of use

For occasional use only (Higdon and Frei 2006; Juliano and Griffiths 2004; Evans and Griffiths 1999)

Risk information

Caution(s) and warning(s)

All products

- ▶ Consult a health care practitioner/health care provider/health care professional/doctor/physician prior to use if you have high blood pressure, glaucoma, and/or overactive bladder syndrome (detrusor instability) (Cornelis and El-Sohemy 2007, Chandrasekaran et al. 2005, Noordzij et al. 2005, Avisar et al. 2002, Arya et al. 2000, Jee et al. 1999, Creighton and Stanton 1990).
- ▶ Avoid taking with health products or foods that contain caffeine and/or increase blood pressure (e.g. medications, coffee, tea, colas, cocoa, guarana, mate, bitter orange extract, synephrine, octopamine, ephedra, ephedrine) (Bui et al. 2006; Bouchard et al. 2005; Haller et al. 2005; FDA 2004; Berardi et al. 2002; Vahedi et al. 2000; Zimmerman 1992; FDA 1988).
- ▶ This product is not intended as a substitute for sleep (Berardi et al. 2002, Zimmerman 1992, FDA 1988).

Products providing 200 mg or more caffeine, per day and recommended for endurance, motor performance, physical energy or to be taken prior to workout

Caffeine has been shown to reduce blood flow to the heart muscle during exercise which might lead to cardiovascular complications such as chest pain, and irregular heartbeat even in healthy individuals. If those symptoms occur, stop use and consult a health care practitioner//health care provider/health care professional/doctor/ physician (Higgins and Babu 2013).

Products providing more than 300 mg caffeine, per day

Consult a health care practitioner/health care provider/health care professional/doctor/physician if you are of childbearing age, pregnant or breastfeeding (Nawrot et al. 2003).

Contraindication(s)

No statement required.

Known adverse reaction(s)

All products

Stop use if hypersensitivity/allergy occurs (Infante et al. 2003; Hinrichs et al. 2002).

Products providing more than 600 mg caffeine, per day

Caffeine may cause anxiety, rapid heart rate (tachycardia), palpitations, insomnia, restlessness, nervousness, tremor and headache (IOM 2001, Zhang 2001, Sawynok 1995).

Non-medicinal ingredients

Must be chosen from the current Natural Health Products Ingredients Database (NHPID) and must meet the limitations outlined in the database.

Storage conditions

No statement required.

Specifications

- ▶ The finished product specifications must be established in accordance with the requirements described in the Natural and Non-prescription Health Products Database (NNHPD) Quality of Natural Health Products Guide.
- ▶ The medicinal ingredient must comply with the requirements outlined in the NHPID.

References cited

Arya LA, Myers DL, Jackson ND. Dietary caffeine intake and the risk for detrusor instability: a case-control study. *Obstetrics and Gynecology* 2000;96(1):85-89.

Ashihara H, Suzuki T. Distribution and biosynthesis of caffeine in plants. *Frontiers in Bioscience* 2004;9:1864-1876.

Avisar R, Avisar E, Weinberger D. Effect of coffee consumption on intraocular pressure. *The Annals of Pharmacotherapy* 2002;36(6):992-995.

Berardi RR, DeSimone EM, Newton GD, Oszko MA, Popovich NG, Rollins CJ, Shimp LA, Tietze KJ, editors. Handbook of Nonprescription Drugs: An Interactive Approach to Self-Care, 13th edition. Washington (DC): American Pharmaceutical Association; 2002.

Bouchard NC, Howland MA, Greller HA, Hoffman RS, Nelson LS. Ischemic stroke associated with use of an ephedra-free dietary supplement containing synephrine. *Mayo Clinic Proceedings* 2005;80(4):541-545.

Bui LT, Nguyen DT, Ambrose PJ. Blood pressure and heart rate effects following a single dose of bitter orange. *The Annals of Pharmacotherapy* 2006;40(1):53-57.

Chandrasekaran S, Rohtchina E, Mitchell P. Effects of caffeine on intraocular pressure: the Blue Mountains Eye Study. *Journal of Glaucoma* 2005;14(6):504-507.

Christopher G, Sutherland D, Smith A. Effects of caffeine in non-withdrawn volunteers. *Human Psychopharmacology* 2005;20(1):47-53.

Cornelis MC, El-Sohehy A. Coffee, caffeine, and coronary heart disease. *Current Opinion in Lipidology* 2007;18(1):13-19.

Creighton SM, Stanton SL. Caffeine: does it affect your bladder? *British Journal of Urology* 1990;66(6):613-614.

Doherty M, Smith PM. Effects of caffeine ingestion on rating of perceived exertion during and after exercise: a meta-analysis. *Scandinavian Journal of Medicine & Science in Sports* 2005;15(2):69-78.

Evans SM, Griffiths RR. Caffeine withdrawal: a parametric analysis of caffeine dosing conditions. *The Journal of Pharmacology and Experimental Therapeutics* 1999;289(1):285-294.

FDA 2004: Food and Drug Administration. Sales of supplements containing ephedrine alkaloids (ephedra) prohibited. Washington (DC): Food and Drug Administration, US Department of Health and Human Services; 2004. [Accessed 2018 June 15]. Available from: <http://www.fda.gov/oc/initiatives/ephedra/february2004/>.

FDA 1988: Food and Drug Administration. 21 CFR Part 340. Stimulant drug products for over-the-counter human use; final monograph; final rule. Washington (DC): U.S. Food and Drug Administration, Department of Health and Human Services; 1988. [Accessed 2018 June 15]. Available from: <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfcfr/CFRSearch.cfm?CFRPart=340&showFR=1>

Gennaro AR, editor. Remington: The Science and Practice of Pharmacy, 20th edition. Washington (DC): Lippincott Williams and Wilkins; 2000.

Greden JF. Anxiety or caffeinism: a diagnostic dilemma. *American Journal of Psychiatry* 1974;131(10):1089-1092.

Haller CA, Benowitz NL, Jacob P. Hemodynamic effects of ephedra-free weight-loss supplements in humans. *The American Journal of Medicine* 2005;118(9):998-1003.

Higdon JV, Frei B. Coffee and health: a review of recent human research. *Critical Reviews in Food Science and Nutrition* 2006;46(2):101-123.

Higgins JP, Babu KM. Caffeine reduces myocardial blood flow during exercise. *The American Journal of Medicine* 2013;126(8):730-e 1-8.

Hinrichs R, Hunzelmann N, Ritzkowsky A, Zollner TM, Krieg T, Scharffetter-Kochanek K. Caffeine hypersensitivity. *Allergy* 2002;57(9):859-60.

Infante S, Baeza ML, Calvo M, De Barrio M, Rubio M, Herrero T. Anaphylaxis due to caffeine. *Allergy* 2003;58(7):681-682.

IOM 2003: Institute of Medicine. Committee on Food Chemicals Codex, Food and Nutrition Board, Institute of Medicine. *Food Chemicals Codex*, 5th edition. Washington (DC): National Academies Press; 2003.

IOM 2001: Institute of Medicine. Committee on Military Nutrition Research, Food and Nutrition Board, Institute of Medicine. *Caffeine for the Sustainment of Mental Task Performance: Formulations for Military Operations*. Washington (DC): National Academy Press; 2001 [Accessed 2018 June 15]. Available from: <http://books.nap.edu/openbook.php?isbn=0309082587>.

Jee SH, He J, Whelton PK, Suh II, Klag MJ. The effect of chronic coffee drinking on blood pressure: a meta-analysis of controlled clinical trials. *Hypertension* 1999;33(2):647-652.

Juliano LM, Griffiths RR. A critical review of caffeine withdrawal: empirical validation of symptoms and signs, incidence, severity, and associated features. *Psychopharmacology* 2004;176(1):1-29.

Kamimori GH, Penetar DM, Headley DB, Thorne DR, Otterstetter R, Belenky G. Effect of three caffeine doses on plasma catecholamines and alertness during prolonged wakefulness. *European Journal of Clinical Pharmacology* 2000;56(8):537-544.

Nawrot P, Jordan S, Eastwood J, Rotstein J, Hugenholtz A, Feeley M. Effects of caffeine on human health. *Food Additives and Contaminants* 2003;20(1):1-30.

Neuhäuser-Berthold M, Beine S, Verwied SC, Lührmann PM. Coffee consumption and total body water homeostasis as measured by fluid balance and bioelectrical impedance analysis. *Annals of Nutrition and Metabolism* 1997;41(1):29-36.

Noordzij M, Uiterwaal CS, Arends LR, Kok FJ, Grobbee DE, Geleijnse JM. Blood pressure response to chronic intake of coffee and caffeine: a meta-analysis of randomized controlled trials. *Journal of Hypertension* 2005;23(5):921-928.

Philip P, Taillard J, Moore N, Delord S, Valtat C, Sagaspe P, Bioulac B. The effects of coffee and napping on night time highway driving: a randomized trial. *Annals of Internal Medicine* 2006;144(11):785-791.

Sawynok J. Pharmacological rationale for the clinical use of caffeine. *Drugs* 1995;49(1):37-50.

Shirley DG, Walter SJ, Noormohamed FH. Natriuretic effect of caffeine: assessment of segmental sodium reabsorption in humans. *Clinical Science* 2002;103(5):461-466.

Smith A, Sutherland D, Christopher G. Effects of repeated doses of caffeine on mood and performance of alert and fatigued volunteers. *Journal of Psychopharmacology* 2005;19(6):620-626.

USP 30 2007: United States Pharmacopeia and the National Formulary (USP 30 - NF 25). Rockville (MD): United States Pharmacopeial Convention, Inc.; 2007.

Vahedi K, Domingo V, Amarenco P, Bousser MG. Ischaemic stroke in a sportsman who consumed MaHuang extract and creatine monohydrate for body building. *Journal of Neurology, Neurosurgery and Psychiatry* 2000;68(1):112-113.

Zajac MA, Zakrzewski AG, Kowal MG, Narayan S. A novel method of caffeine synthesis from uracil. *Synthetic Communications* 2003;33(19):3291-3297.

Zhang WY. A benefit-risk assessment of caffeine as an analgesic adjuvant. *Drug Safety* 2001;24(15):1127-1142.

Zimmerman DR. Zimmerman's Complete Guide to Nonprescription Drugs, 2nd edition. Detroit (MI): Gale Research Inc.; 1992.

Zwyghuizen-Doorenbos A, Roehrs TA, Lipschutz L, Timms V, Roth T. Effects of caffeine on alertness. *Psychopharmacology* 1990;100(1):36-39.

References reviewed

- Ahmed SV, Jayawarna C, Jude E. Post lumbar puncture headache: diagnosis and management. *Postgraduate Medical Journal* 2006;82(973):713-716.
- Ahrens JN, Crixell SH, Lioyd LK, Walker JL. The physiological effects of caffeine in women during treadmill walking. *Journal of Strength and Conditioning Research* 2007;21(1):164-168.
- Ali M, Afzal M. A potent inhibitor of thrombin stimulated platelet thromboxane formation from unprocessed tea. *Prostaglandins, Leukotrienes and Medicine* 1987;27(1):9-13.
- Aqel RA, Zoghbi GJ, Trimm JR, Baldwin SA, Iskandrian AE. Effect of caffeine administered intravenously on intracoronary-administered adenosine-induced coronary hemodynamics in patients with coronary artery disease. *The American Journal of Cardiology* 2004;93(3):343-346.
- Ascherio A, Weisskopf MG, O'Reilly EJ, McCullough ML, Calle EE, Rodriguez C, Thun MJ. Coffee consumption, gender, and Parkinson's disease mortality in the cancer prevention study II cohort: the modifying effects of estrogen. *American Journal of Epidemiology* 2004;160(10):977-984.
- Astrup A, Toubro S, Cannon S, Hein P, Breum L, Madsen J. Caffeine: a double-blind, placebo controlled study of its thermogenic, metabolic, and cardiovascular effects in healthy volunteers. *The American Journal of Clinical Nutrition* 1990;51(5):759-767.
- Azcona O, Barbanoj MJ, Torrent J, Jane F. Evaluation of the central effects of alcohol and caffeine interaction. *British Journal of Clinical Pharmacology* 1995;40(4):393-400.
- Balat O, Balat A, Ugur MG, Pence S. The effect of smoking and caffeine on the fetus and placenta in pregnancy. *Clinical and Experimental Obstetrics and Gynecology* 2003;30(1):57-59.
- Bara AI, Barley EA. Caffeine for asthma. Oxfordshire (UK): The Cochrane Database of Systematic Reviews; 2001, Issue 4. Art. No.: CD001112. DOI: 10.1002/14651858.CD001112.
- Barnett G, Segura J, De La Torre R, Carbo M. Pharmacokinetic determination of relative potency of quinolone inhibition of caffeine disposition. *European Journal of Clinical Pharmacology* 1990;39(1):63-69.
- Barone, JJ, Roberts HR. Caffeine consumption. *Food and Chemical Toxicology* 1996;34(1):119-129.
- Beach CA, Mays DC, Guiler RC, Jacober CH, Gerber N. Inhibition of elimination of caffeine by disulfiram in normal subjects and recovering alcoholics. *Clinical Pharmacology and Therapeutics* 1986;39(3):265-270.

Bell DG, Jacobs I, Ellerington K. Effect of caffeine and ephedrine ingestion on anaerobic exercise performance. *Medicine and Science in Sports and Exercise* 2001;33(8):1399-1403.

Bell DG, McLellan TM, Sabiston CM. Effect of ingesting caffeine and ephedrine on 10-km run performance. *Medicine and Science in Sports and Exercise* 2002;34(2):344-349.

Belza A, Frandsen E, Kondrup J. Body fat loss achieved by stimulation of thermogenesis by a combination of bioactive food ingredients: a placebo-controlled, double-blind 8-week intervention in obese subjects. *International Journal of Obesity* 2007;31(1):121-130.

Bernstein GA, Carroll ME, Crosby RD, Perwien AR, Go FS, Benowitz NL. Caffeine effects on learning, performance, and anxiety in normal school-age children. *Journal of the American Academy of Child and Adolescent Psychiatry* 1994;33(3):407-415.

Bott Kaspar. 1983. Preparation of caffeine. United States Patent US 4380631. 1983-04-19.

Brown NJ, Ryder D, Branch RA. A pharmacodynamic interaction between caffeine and phenylpropanolamine. *Clinical Pharmacology and Therapeutics* 1991;50(4):363-371.

Bruce CR, Anderson ME, Fraser SF, Stepto NK, Klein R, Hopkins WG, Hawley JA. Enhancement of 2000-m rowing performance after caffeine ingestion. *Medicine and Science in Sports and Exercise* 2000;32(11):1958-1963.

Carbo M, Segura J, De La Torre R, Badenas JM, Cami J. Effect of quinolones on caffeine disposition. *Clinical Pharmacology and Therapeutics* 1989;45(3):234-240.

Carrillo JA, Benitez J. Clinically significant pharmacokinetic interactions between dietary caffeine and medications. *Clinical Pharmacokinetics* 2000;39(2):127-153.

Cavalcante JW, Santos PR, Menezes MG, Marques HO, Cavalcante LP, Pacheco WS. Influence of caffeine on blood pressure and platelet aggregation. *Arquivos Brasileiros de Cardiologia* 2000;75(2):97-105.

Choi A, Laurito CE, Cunningham FE. Pharmacologic management of postdural puncture headache. *Annals of Pharmacotherapy* 1996;30(7-8):831-839.

Culm-Merdek KE, Von Moltke LL, Harmatz JS, Greenblatt DJ. Fluvoxamine impairs single-dose caffeine clearance without altering caffeine pharmacodynamics. *British Journal of Clinical Pharmacology* 2005;60(5):486-493.

Devoe LD, Murray C, Youssif A, Arnaud M. Maternal caffeine consumption and fetal behaviour in the normal third-trimester pregnancy. *American Journal of Obstetrics and Gynecology* 1993;168(4):1105-1111.

Dickerson LM, Mazyck PJ, Hunter MH. Premenstrual syndrome. *American Family Physician* 2003;67(8):1743-1752.

Du Y, Melchert HU, Knopf H, Braemer-Hauth M, Gerding B, Pabel E. Association of serum caffeine concentrations with blood lipids in caffeine-drug users and nonusers - results of German National Health Surveys from 1984 to 1999. *European Journal of Epidemiology* 2005;20(4):311-316.

Edwards RG. An evaluation of the health aspects of caffeine as a food ingredient. Washington (DC): Arent Fox Kintner Plotkin & Kahn; 1997.

FDA 1988a: Food and Drug Administration. Federal Register Volume 53, Number 39, page 6105. Washington (DC): U.S. Food and Drug Administration, Department of Health and Human Services. February 29, 1988.

FDA 1988b: Food and Drug Administration. Federal Register Volume 53, Number 221, page 46201. Washington (DC): U.S. Food and Drug Administration, Department of Health and Human Services. November 16, 1988.

Foxx RM, Rubinoff A. Behavioral treatment of caffeinism: reducing excessive coffee drinking. *Journal of Applied Behavior Analysis* 1979;12(3):335-344.

Fudin R, Nicastro R. Can caffeine antagonize alcohol-induced performance decrements in humans? *Perceptual and Motor Skills* 1988;67(2):375-391.

Goldstein J, Silberstein SD, Saper JR, Ryan RE, Lipton RB. Acetaminophen, aspirin, and caffeine in combination versus ibuprofen for acute migraine: results from a multicenter, double blind, randomized, parallel-group, single-dose, placebo-controlled study. *Headache* 2006;46(3):444-453.

Greenberg JA, Boozer CN, Geliebter A. Coffee, diabetes, and weight control. *The American Journal of Clinical Nutrition* 2006;84(4):682-693.

Grobbee DE, Rimm EB, Giovannucci E, Colditz G, Stampfer M, Willett W. Coffee, caffeine, and cardiovascular disease in men. *New England Journal of Medicine* 1990;323(15):1026-1032.

Hagg S, Spigset O, Mjorndal T, Dahlqvist R. Effect of caffeine on clozapine pharmacokinetics in healthy volunteers. *British Journal of Clinical Pharmacology* 2000;49(1):59-63.

Hallstorm H, Wolk A, Gynn A, Michaelsson K. Coffee, tea and caffeine consumption in relation to osteoporotic fracture risk in a cohort of Swedish women. *Osteoporosis International*. 2006;17(7):1055-1064.

Haskell CF, Kennedy DO, Wesnes KA, Milne AL, Scholey AB. Double-blind, placebo controlled, multi-dose evaluation of the acute behavioural effects of guarana in humans. *Journal of Psychopharmacology* 2007;21(1):65-70.

Ishizuk H, Eguchi H, Oda T, Ogawaf S, Nakagawa K, Honjo S, Kono S. Relation of coffee, green tea, and caffeine intake to gallstone disease in middle-aged Japanese men. *European Journal of Epidemiology* 2003;18(5):401-405.

Jeppesen U, Loft S, Poulsen HE, Brsen K. A fluvoxamine-caffeine interaction study. *Pharmacogenetics* 1996;6(3):213-222.

Joeres R, Klinker H, Heusler H, Epping J, Richter E. Influence of mexiletine on caffeine elimination. *Pharmacology and Therapeutics* 1987;33(1):163-169.

Kiyohara C, Kono S, HonJo S, Todoroki I, Sakurai Y, Nishikawa H, Kogo H, Ogawa S, Nakagawa K. Inverse association between coffee drinking and serum uric acid concentrations in middle-aged Japanese males. *British Journal of Nutrition* 1999;82(2):125-130.

Kleemola P, Jousilahti P, Pietinen P, Vartiainen E, Tuomilehto J. Coffee consumption and the risk of coronary heart disease and death. *Archives of Internal Medicine* 2000;160(22):3393-4000.

Labbe L, Abolfathi Z, Robitaille NM, St-Maurice F, Gilbert M, Turgeon J. Stereoselective disposition of the antiarrhythmic agent mexiletine during the concomitant administration of caffeine. *Therapeutic Drug Monitoring* 1999;21(2):191-199.

Lake CR, Rosenberg DB, Gallant S, Zaloga G, Chernow B. Phenylpropanolamine increases plasma caffeine levels. *Clinical Pharmacology and Therapeutics* 1990;47(6):675-685.

Laska EM, Sunshine A, Mueller F, Elvers WB, Siegel C, Rubin A. Caffeine as an analgesic adjuvant. *Journal of the American Medical Association* 1984;251(130):1711-1718.

Leitzmann MF, Stampfer MJ, Willett WC, Spiegelman D, Colditz GA, Giovannucci EL. Coffee intake is associated with lower risk of symptomatic gallstone disease in women. *Gastroenterology* 2002;123(6):1823-1830.

Leitzmann MF, Willett WC, Rimm EB, Stampfer MJ, Spiegelman D, Colditz GA, Giovannucci E. A prospective study of coffee consumption and the risk of symptomatic gallstone disease in men. *The Journal of the American Medical Association* 1999;281(22):2106-2112.

Leitzmann MF, Willett WC, Rimm EB, Stampfer MJ, Spiegelman D, Colditz GA, Giovannucci E, Lin W, Geiderman J. Myth: fluids, bed rest, and caffeine are effective in preventing and treating patients with post-lumbar puncture headache. *The Western Journal of Medicine* 2002;176(1):69-70.

Lipton RB, Stewart WF, Ryan RE, Saper J, Silberstein S, Sheftell F. Efficacy and safety of acetaminophen, aspirin, and caffeine in alleviating migraine headache pain: three double-blind, randomized, placebo-controlled trials. *Archives of Neurology* 1998;55(2):210-217.

Maillard C, Babadjamian A, Balansard G, Ollivier B, Bamba D. Study of caffeine - catechin association in lyophilized fresh seeds and in stabilized extract of *cola nitida*. *Planta Medica* 1985;51(6):515-517.

Malinow MR, Bostom AG, Krauss RM. Homocyst(e)ine, diet, and cardiovascular diseases: a statement for healthcare professionals from the Nutrition Committee, American Heart Association. *Circulation* 1999;99(1):178-182.

May DC, Jarboe CH, VanBakel AB, Williams WM. Effects of cimetidine on caffeine disposition in smokers and non-smokers. *Clinical Pharmacology and Therapeutics* 1982;31(5):656-661.

Meyers BM, Cafarelli E. Caffeine increases time to fatigue by maintaining force and not by altering firing rates during submaximal isometric contractions. *Journal of Applied Physiology* 2005;99(3):1056-1063.

Migliardi JR, Armellino JJ, Friedman M, Gillings DB, Beaver WT. Caffeine as an analgesic adjuvant in tension headache. *Clinical Pharmacology and Therapeutics* 1994;56(5):576-586.

Min B, Cios D, Kluger J, White CM. Absence of QTc-interval-prolonging or hemodynamic effects of a single dose of bitter-orange extract in healthy subjects. *Pharmacotherapy* 2005;25(12):1719-1724.

Myers MG, Basinski A. Coffee and coronary heart disease. *Archives of Internal Medicine* 1992;152(9):1767-1772.

Norlock FE. Benign breast pain in women: a practical approach to evaluation and treatment. *Journal of the American Medical Women's Association* 2002;57(2):85-90.

Olak-Bialon B, Marcisz C, Jonderko G, Olak Z, Szymyszal J, Orzel A. [Does coffee drinking influence serum uric acid concentration?] *Wiadomości lekarskie* 2004;57(Supplement 1):233-237 (in Polish).

Papaioannou TG, Vlachopoulos C, Ioakeimidis N, Alexopoulos N, Stefanadis C. Nonlinear dynamics of blood pressure variability after caffeine consumption. *Clinical Medicine and Research* 2006;4(2):114-118.

Passmore AP, Kondowe GB, Johnston GD. Renal and cardiovascular effects of caffeine: a doseresponse study. *Clinical Science* 1987;72(6):749-756.

Phillips-Bute BG, Lane JD. Caffeine withdrawal symptoms following brief caffeine deprivation. *Physiological Behavior* 1997;63(1):35-39.

Pollock BG, Wylie M, Stack JA, Sorisio DA, Thompson DS, Kirshner MA, Folan MM, Condifer KA. Inhibition of caffeine metabolism by estrogen replacement therapy in postmenopausal women. *Journal of Clinical Pharmacology* 1999;39(9):936-940.

Przybilla B, Ring J, Burg G. [Anaphylaxis following ingestion of coffee, chronic urticaria and analgesics idiosyncrasy]. *Der Hautarzt; Zeitschrift für Dermatologie, Venerologie, und verwandte Gebiete* 1983;34(2):73-76 (in German).

Raaska K, Raitasuo V, Laitila J, Neuvonen PJ. Effect of caffeine-containing versus decaffeinated coffee on serum clozapine concentrations in hospitalised patients. *Basic and Clinical Pharmacology and Toxicology* 2004;94(1):13-18.

Rapuri PB, Gallagher JC, Kinyamu HK, Ryschon KL. Caffeine intake increases the rate of bone loss in elderly women and interacts with vitamin D receptor genotypes. *American Journal of Clinical Nutrition* 2001;74(5):694-700.

Rasheed P, Al-Sowielem LS. Prevalence and predictors of premenstrual syndrome among college-aged women in Saudi Arabia. *Annals of Saudi Medicine* 2003;23(6):381-387.

Riesenhuber A, Boehm M, Posch M, Aufricht C. Diuretic potential of energy drinks. *Amino Acids* 2006;31(1):81-83.

Ross GW, Abbott RD, Petrovitch H, Morens DM, Grandinetti A, Tung KH, Tanner CM, Masaki KH, Blanchette PL, Curb JD, Popper JS, White LR. Association of coffee and caffeine intake with the risk of Parkinson's disease. *The Journal of the American Medical Association* 2000;283(20):2674-2679.

Rossignol AM. Caffeine-containing beverages and premenstrual syndrome in young women. *American Journal of Public Health* 1985;75(11):1335-1337.

Rossignol AM, Bonnlander H. Caffeine-containing beverages, total fluid consumption, and premenstrual syndrome. *American Journal of Public Health* 1990;80(9):1106-1110.

Rossignol AM, Bonnlander H, Song L, Phillis JW. Do women with premenstrual symptoms self-medicate with caffeine? *Epidemiology* 1991;2(6):403-408.

Rossignol AM, Zhang JY, Chen YZ, Xiang Z. Tea and premenstrual syndrome in the People's Republic of China. *American Journal of Public Health* 1989;79(1):67-69.

Salazar-Martinez E, Willett WC, Ascherio A, Manson JE, Leitzmann MF, Stampfer MJ, Hu FB. Coffee consumption and risk for type 2 diabetes mellitus. *Annals of Internal Medicine* 2004;140(1):1-8.

Sanderink GJ, Bournique B, Stevens J, Petry M, Martinet M. Involvement of human CYP1A isoenzymes in the metabolism and drug interactions of riluzole in vitro. *The Journal of Pharmacology and Experimental Therapy* 1997;282(3):1465-1472.

Santos IS. Caffeine intake and pregnancy outcomes: a meta-analytic review. *Cadernos de Saúde Pública* 1998;14(3):523-530.

Schmidt B, Roberts RS, Davis P, Doyle LW, Barrington KJ, Ohlsson A, Solimano A, Tin W. Caffeine therapy for apnea of prematurity. *The New England Journal of Medicine* 2006;354(20):2112-2121.

Simon O, Cohen L, Pierrot-Deseilligny C. [Post-lumbar puncture syndrome. Causes, prevention, treatments]. *Revue Neurologique* 1997;153(10):605-608 (in French).

Smith A. Effects of caffeine on human behavior. *Food and Chemical Technology* 2002;40(9):1243-1255.

Smith B, Wingard DL, Smith TC, Kritz-Silverstein D, Barrett-Connor E. Does coffee consumption reduce the risk of type 2 diabetes in individuals with impaired glucose? *Diabetes Care* 2006;29(11):2385-2390.

Smith PF, Smith A, Miners J, McNeil J, Proudfoot A. Report from the Expert Working Group on the Safety Aspects of Dietary Caffeine. Canberra (AU): Australian New Zealand Food Authority; 2000.

Stavric B. An update on research with coffee/caffeine (1989-1990). *Food and Chemical Toxicology* 1992;30(6):533-555.

Steer PA, Flenady VJ, Shearman A, Lee TC, Tudehope DI, Charles BG. Periextubation caffeine in preterm neonates: a randomized dose response trial. *Journal of Paediatrics and Child Health* 2003;39(7):511-515.

Stuart GR, Hopkin WG, Cook C, Cairns SP. Multiple effects of caffeine on simulated highintensity team-sport performance. *Medicine and Science in Sports and Exercise* 2005;37(11):1998-2005.

Urgert R, Katan MB. The cholesterol-raising factor from coffee beans. *Annual Review of Nutrition* 1997;17:305-324.

Urgert R, Van Vliet T, Zock PL, Katan MB. Heavy coffee consumption and plasma homocysteine: a randomized controlled trial in healthy volunteers. *The American Journal of Clinical Nutrition* 2000;72(5):1107-1110.

Van Dusseldorp M, Katan MB. Headache caused by caffeine withdrawal among moderate coffee drinkers switched from ordinary to decaffeinated coffee: a 12 week double blind trial. *British Medical Journal* 1990;300(6739):1558-1559.

Vandenbergh K, Gillis N, Van Leemputte M, Van Hecke P, Vanstapel F, Hespel P. Caffeine counteracts the ergogenic action of muscle creatine loading. *Journal of Applied Physiology* 1996;80(2):452-457.

Varani K, Portaluppi F, Gessi S, Merighi S, Ongini E, Belardinelli L, Borea PA. Dose and time effects of caffeine intake on human platelet adenosine A_{2A} receptors: functional and biochemical aspects. *Circulation* 2000;102(3):285-289.

Verhoef P, Pasman WJ, Van Vliet T, Urgert R, Katan MB. Contribution of caffeine to the homocysteine-raising effect of coffee: a randomized controlled trial in humans. *The American Journal of Clinical Nutrition* 2002;76(6):1244-1248.

Wahlander A, Paumgartner G. Effect of ketoconazole and terbinafine on the pharmacokinetics of caffeine in healthy volunteers. *European Journal of Clinical Pharmacology* 1989;37(3):279-283.

Watson JM, Jenkins EJ, Hamilton P, Lunt MJ, Kerr D. Influence of caffeine on the frequency and perception of hypoglycemia in free-living patients with type 1 diabetes. *Diabetes Care* 2000;23(4):455-459.

Westertep-Planteqa MS, Ljeune MP, Kovacs EM. Body weight loss and weight maintenance in relation to habitual caffeine intake and green tea supplementation. *Obesity Research* 2005;13(7):1195-1204.

Yucel A, Ozyalcin S, Talu GK, Yucel EC, Erdine S. Intravenous administration of caffeine sodium benzoate for postdural puncture headache. *Regional Anesthesia and Pain Medicine* 1999;24(1):51-54.